

# SUBSTITUTE SPECIFICATION

## DNA ENCODING A GROWTH FACTOR SPECIFIC FOR EPITHELIAL CELLS

5        This application is a divisional of application  
Serial No. 08/106,775, filed August 16, 1993, which is a  
continuation of Serial No. 07/780,845, filed October 23,  
1991, which is a continuation of Serial No. 07/304,281,  
filed January 31, 1989.

### FIELD OF THE INVENTION

10        The present invention relates to growth factors,  
particularly to isolation of a polypeptide growth factor  
similar to a family of factors including known fibroblast  
growth factors (FGFs). This invention also relates to  
construction of complementary DNA (cDNA) segments from  
15        messenger RNA (mRNA) encoding the novel growth factor.  
Further, this invention pertains to synthesis of products  
of such DNA segments by recombinant cells, and to the  
manufacture and use of certain other novel products  
enabled by the identification and cloning of DNAs  
20        encoding this growth factor.

ABBREVIATIONS USED IN THIS APPLICATION

	aFGF	acidic fibroblast growth factor
	bFGF	basic fibroblast growth factor
	EGF	epidermal growth factor
5	HSAC	heparin-Sepharose affinity chromatography
	kb	kilobases
	kDa	kilodaltons
	KGF	keratinocyte growth factor
10	NaDodSO <sub>4</sub> /PAGE	Sodium dodecylsulphate (SDS)/polyacryl- amide gel electrophoresis
	RP-HPLC	reversed-phase high performance liquid chromatography
	TGF $\alpha$	transforming growth factor $\alpha$

# BACKGROUND OF THE INVENTION

Growth factors are important mediators of inter-cellular communication. These potent molecules are generally released by one cell type and act to influence proliferation of other cell types (James, R. and Bradshaw, R.A. (1984), *Ann. Rev. Biochem.* 53, 259-292). Interest in growth factors has been heightened by evidence of their potential involvement in neoplasia (Sporn, M. B. and Todaro, G. J. (1980), *N. Eng. J. Med.* 303, 878-880). The v-sis transforming gene of simian sarcoma virus encodes a protein that is homologous to the B chain of platelet-derived growth factor (James, R. and Bradshaw, R.A. (1984) *Ann. Rev. Biochem.* 53, 259-292; Doolittle, R. F., et al. (1983) *Science* 221, 275-277). Moreover, a number of oncogenes are homologues of genes encoding growth factor receptors (James, R. and Bradshaw, R.A. (1984) *Ann. Rev. Biochem.* 53, 259-292). Thus, increased understanding of growth factors and their receptor-mediated signal transduction pathways is likely to provide insights into mechanisms of both normal and malignant cell growth.

One known family of growth factors affecting connective tissue cells includes acidic fibroblast growth factor (aFGF), basic fibroblast growth factor (bFGF), and the related products of the *hst*, FGF-5 and *int-2* oncogenes.

Further, it is known that some growth factors, including the following, have heparin-binding properties: aFGF (Maciag, T., Mehlman, T., Friesel, R. and Schreiber, A.B. (1984) *Science* 225, 932-935; Conn, G. and Hatcher, V.B. (1984) *Biochem. Biophys. Res. Comm.* 124, 262-268); bFGF (Gospodarowicz, D., Cheng, J., Lui, G.-M., Baird, A. and Bohlen, P. (1984) *Proc. Natl. Acad. Sci. USA* 81, 6963-6967; Maciag, T., Mehlman, T., Friesel, R. and Schreiber, A.B. (1984) *Science* 225, 932-935); granulocyte/macrophage colony stimulating factor (James, R. and Bradshaw, R.A. (1984) *Ann. Rev. Biochem.* 53, 259-292); and interleukin 3 (James, R. and Bradshaw, R.A. (1984)

Ann. Rev. Biochem. 53, 259-292). Each of these polypeptide factors is produced by stromal cells (James, R. and Bradshaw, R.A. (1984) Ann. Rev. Biochem. 53, 259-292, Doolittle, R.F., Hunkapiller, M.W., Hood, L.E., Devare, S.G., Robbins, K.C., Aaronson, S.A. and Antoniades, M.N. (1983) Science 221, 275-277, Roberts, R., Gallagher, J., Spooncer, E., Allen, T.D., Bloomfield, F. and Dexter, T. M. (1988) Nature 332, 376-378). Such factors appear to be deposited in the extracellular matrix, or on proteoglycans coating the stromal cell surface (James, R. and Bradshaw, R.A. (1984) Ann. Rev. Biochem. 53, 259-292, Roberts, R., Gallagher, J., Spooncer, E., Allen, T.D., Bloomfield, F. and Dexter, T. M. (1988) Nature 332, 376-378). It has been postulated that their storage, release and contact with specific target cells are regulated by this interaction (Roberts, R., Gallagher, J., Spooncer, E., Allen, T.D., Bloomfield, F. and Dexter, T. M. (1988) Nature 332, 376-378, Vlodavsky, I., Folkman, J., Sullivan, R., Fridman, R., Ishai-Michaeli, R., Sasse, J. and Klagsburn, M. (1987) Proc. Natl. Acad. Sci. USA 84, 2292-2296).

It is widely recognized, however, that the vast majority of human malignancies are derived from epithelial tissues (Wright, N. and Allison, M. (1984) The Biology of Epithelial Cell Populations (Oxford University Press, New York) Vol. 1, pp. 3-5). Effectors of epithelial cell proliferation derived from mesenchymal cells have been described (James, R. and Bradshaw, R.A. (1984) Ann. Rev. Biochem. 53, 259-292, Doolittle, R.F., Hunkapiller, M.W., Hood, L.E., Devare, S.G., Robbins, K.C., Aaronson, S.A. and Antoniades, M.N. (1983) Science 221, 275-277, Waterfield, M.D., Scrace, G.J., Whittle, N., Strooband, P., Johnson, A., Wasteton, A., Westermarck, B., Heldin, C.-H., Huang, J.S. and Deuel, T.F. (1983) Nature 304, 35-39), however, their molecular identities and structures have not been elucidated.

In light of this dearth of knowledge about such mesenchymal growth factors affecting epithelial cells, it is apparent that there has been a need for methods and

compositions and bioassays which would provide an improved knowledge and analysis of mechanisms of regulation of epithelial cell proliferation, and, ultimately, a need for novel diagnostics and therapies based on the factors involved therein.

This invention contemplates the application of methods of protein isolation and recombinant DNA technologies to fulfill such needs and to develop means for producing protein factors of mesenchymal origin, which appear to be related to epithelial cell proliferation processes and which could not be produced otherwise. This invention also contemplates the application of the molecular mechanisms of these factors related to epithelial cell growth processes.

#### SUMMARY OF THE INVENTION

The present invention relates to developments of protein isolation and recombinant DNA technologies, which include production of novel growth factor proteins affecting epithelial cells, free of other peptide factors. Novel DNA segments and bioassay methods are also included.

The present invention in particular relates to a novel protein having structural and/or functional characteristics of a known family of growth factors which includes acidic fibroblast growth factor (aFGF), basic fibroblast growth factor (bFGF) and the related products of the *hst*, FGF-5 and *int-2* oncogenes. This new member of the FGF polypeptide family retains the heparin-binding properties of the FGFs but has evolved a unique target cell specificity. This growth factor appears to be specific for epithelial cells and is particularly active on keratinocytes. Therefore, this novel factor has been designated "keratinocyte growth factor" (KGF). Notwithstanding its lack of activity on fibroblasts, since it is the sixth known member of the FGF polypeptide family, KGF may also be referred to as FGF-6.

Accordingly, this invention relates, in part, to purified KGF or KGF-like proteins and methods for preparing these proteins. Such purified factors may be made by cultivation of human cells which naturally secrete these proteins and application of isolation methods according to the practice of this invention. These proteins can be used for biochemical and biological studies leading, for example, to isolation of DNA segments encoding KGF or KGF-like polypeptides.

The present invention also relates to such DNA segments which encode KGF or KGF-like proteins. In a principal embodiment, the present invention relates to DNA segments, which encode KGF-related products, consisting of: human cDNA clones 32 or 49, derived from polyadenylated RNA extracted from the human embryonic lung fibroblast cell line M426; recombinants and mutants of these clones; and related DNA segments which can be detected by hybridization to any of the above human DNA segments, which related segments encode KGF-like proteins or portions thereof.

In the practice of one embodiment of this invention, the DNA segments of the invention are capable of being expressed in suitable host cells, thereby producing KGF or KGF-like proteins. The invention also relates to m-RNAs produced as the result of transcription of the sense strands of the DNA segments of this invention.

In another embodiment, the invention relates to a recombinant DNA molecule comprising a vector and a DNA of the present invention. These recombinant molecules are exemplified by molecules comprising a KGF cDNA and any of the following vector DNAs: a bacteriophage  $\lambda$  cloning vector (exemplified by  $\lambda$ pCEV9); a DNA sequencing plasmid vector (e.g., a pUC variant); a bacterial gene expression vector (e.g., pKK233-2); or a mammalian gene expression vector (such as pMMT).

In still another embodiment, the invention comprises a cell, preferably a mammalian cell, transformed with a DNA of the invention. Further, the invention comprises cells, including insect cells, yeast cells and bacterial

cells such as those of *Escherichia coli* and *B. subtilis*, transformed with DNAs of the invention. According to another embodiment of this aspect of the invention, the transforming DNA is capable of being expressed in the cell, thereby increasing in the cell the amount of KGF or KGF-like protein encoded by this DNA.

The primary KGF translation product predicted from its cDNA sequence contains an N-terminal hydrophobic region which likely serves as a signal sequence for secretion and which is not present in the mature KGF molecule. In a most preferred embodiment of the gene expression aspect of the invention, the cell transformed by the DNA of the invention secretes the protein encoded by that DNA in the (truncated) form that is secreted by human embryonic lung fibroblast cells.

Still further, this invention contemplates KGF or KGF-like proteins produced by expression of a DNA of the invention, or by translation of an RNA of the invention. Preferably, these proteins will be of the secreted form (i.e., lacking an apparent signal sequence). These protein factors can be used for functional studies, and can be purified for additional structural and functional analyses, such as qualitative and quantitative receptor binding assays.

Moreover, the ability to produce large quantities of this novel growth factor by recombinant techniques will allow testing of its clinical applicability in situations where specific stimulation of growth of epithelial cells is of particular importance. Accordingly, this invention includes pharmaceutical compositions comprising KGF or KGF-like polypeptides for use in the treatment of such conditions, including, for example, healing of wounds due to burns or stimulation of transplanted corneal tissue.

According to this embodiment of the invention, the novel KGF-like proteins will be protein products of "unmodified" DNAs and mRNAs of the invention, or will be modified or genetically engineered protein products. As a result of engineered mutations in the DNA sequences modified KGF-like proteins will have one or more

differences in amino acid sequence from the corresponding naturally occurring "wild-type" proteins. According to one embodiment of this aspect of this invention, the modified KGF-like proteins will include "chimeric" molecules comprising segments of amino acid sequences of KGF and at least one other member of the FGF peptide family.

5  
10  
15  
20  
25  
30  
35  
Ultimately, given results of analogous successful approaches with other peptide factors having similar properties, development of such chimeric KGF-like polypeptides should lead to superior, "second generation" forms of KGF-like peptides for clinical purposes. These modified KGF-like products might be smaller, more stable, more potent, and/or easier or less expensive to produce, for example.

15  
20  
25  
30  
35  
40  
45  
50  
55  
60  
65  
70  
75  
80  
85  
90  
95  
100  
105  
110  
115  
120  
125  
130  
135  
140  
145  
150  
155  
160  
165  
170  
175  
180  
185  
190  
195  
200  
205  
210  
215  
220  
225  
230  
235  
240  
245  
250  
255  
260  
265  
270  
275  
280  
285  
290  
295  
300  
305  
310  
315  
320  
325  
330  
335  
340  
345  
350  
355  
360  
365  
370  
375  
380  
385  
390  
395  
400  
405  
410  
415  
420  
425  
430  
435  
440  
445  
450  
455  
460  
465  
470  
475  
480  
485  
490  
495  
500  
505  
510  
515  
520  
525  
530  
535  
540  
545  
550  
555  
560  
565  
570  
575  
580  
585  
590  
595  
600  
605  
610  
615  
620  
625  
630  
635  
640  
645  
650  
655  
660  
665  
670  
675  
680  
685  
690  
695  
700  
705  
710  
715  
720  
725  
730  
735  
740  
745  
750  
755  
760  
765  
770  
775  
780  
785  
790  
795  
800  
805  
810  
815  
820  
825  
830  
835  
840  
845  
850  
855  
860  
865  
870  
875  
880  
885  
890  
895  
900  
905  
910  
915  
920  
925  
930  
935  
940  
945  
950  
955  
960  
965  
970  
975  
980  
985  
990  
995  
1000  
1005  
1010  
1015  
1020  
1025  
1030  
1035  
1040  
1045  
1050  
1055  
1060  
1065  
1070  
1075  
1080  
1085  
1090  
1095  
1100  
1105  
1110  
1115  
1120  
1125  
1130  
1135  
1140  
1145  
1150  
1155  
1160  
1165  
1170  
1175  
1180  
1185  
1190  
1195  
1200  
1205  
1210  
1215  
1220  
1225  
1230  
1235  
1240  
1245  
1250  
1255  
1260  
1265  
1270  
1275  
1280  
1285  
1290  
1295  
1300  
1305  
1310  
1315  
1320  
1325  
1330  
1335  
1340  
1345  
1350  
1355  
1360  
1365  
1370  
1375  
1380  
1385  
1390  
1395  
1400  
1405  
1410  
1415  
1420  
1425  
1430  
1435  
1440  
1445  
1450  
1455  
1460  
1465  
1470  
1475  
1480  
1485  
1490  
1495  
1500  
1505  
1510  
1515  
1520  
1525  
1530  
1535  
1540  
1545  
1550  
1555  
1560  
1565  
1570  
1575  
1580  
1585  
1590  
1595  
1600  
1605  
1610  
1615  
1620  
1625  
1630  
1635  
1640  
1645  
1650  
1655  
1660  
1665  
1670  
1675  
1680  
1685  
1690  
1695  
1700  
1705  
1710  
1715  
1720  
1725  
1730  
1735  
1740  
1745  
1750  
1755  
1760  
1765  
1770  
1775  
1780  
1785  
1790  
1795  
1800  
1805  
1810  
1815  
1820  
1825  
1830  
1835  
1840  
1845  
1850  
1855  
1860  
1865  
1870  
1875  
1880  
1885  
1890  
1895  
1900  
1905  
1910  
1915  
1920  
1925  
1930  
1935  
1940  
1945  
1950  
1955  
1960  
1965  
1970  
1975  
1980  
1985  
1990  
1995  
2000  
2005  
2010  
2015  
2020  
2025  
2030  
2035  
2040  
2045  
2050  
2055  
2060  
2065  
2070  
2075  
2080  
2085  
2090  
2095  
2100  
2105  
2110  
2115  
2120  
2125  
2130  
2135  
2140  
2145  
2150  
2155  
2160  
2165  
2170  
2175  
2180  
2185  
2190  
2195  
2200  
2205  
2210  
2215  
2220  
2225  
2230  
2235  
2240  
2245  
2250  
2255  
2260  
2265  
2270  
2275  
2280  
2285  
2290  
2295  
2300  
2305  
2310  
2315  
2320  
2325  
2330  
2335  
2340  
2345  
2350  
2355  
2360  
2365  
2370  
2375  
2380  
2385  
2390  
2395  
2400  
2405  
2410  
2415  
2420  
2425  
2430  
2435  
2440  
2445  
2450  
2455  
2460  
2465  
2470  
2475  
2480  
2485  
2490  
2495  
2500  
2505  
2510  
2515  
2520  
2525  
2530  
2535  
2540  
2545  
2550  
2555  
2560  
2565  
2570  
2575  
2580  
2585  
2590  
2595  
2600  
2605  
2610  
2615  
2620  
2625  
2630  
2635  
2640  
2645  
2650  
2655  
2660  
2665  
2670  
2675  
2680  
2685  
2690  
2695  
2700  
2705  
2710  
2715  
2720  
2725  
2730  
2735  
2740  
2745  
2750  
2755  
2760  
2765  
2770  
2775  
2780  
2785  
2790  
2795  
2800  
2805  
2810  
2815  
2820  
2825  
2830  
2835  
2840  
2845  
2850  
2855  
2860  
2865  
2870  
2875  
2880  
2885  
2890  
2895  
2900  
2905  
2910  
2915  
2920  
2925  
2930  
2935  
2940  
2945  
2950  
2955  
2960  
2965  
2970  
2975  
2980  
2985  
2990  
2995  
3000  
3005  
3010  
3015  
3020  
3025  
3030  
3035  
3040  
3045  
3050  
3055  
3060  
3065  
3070  
3075  
3080  
3085  
3090  
3095  
3100  
3105  
3110  
3115  
3120  
3125  
3130  
3135  
3140  
3145  
3150  
3155  
3160  
3165  
3170  
3175  
3180  
3185  
3190  
3195  
3200  
3205  
3210  
3215  
3220  
3225  
3230  
3235  
3240  
3245  
3250  
3255  
3260  
3265  
3270  
3275  
3280  
3285  
3290  
3295  
3300  
3305  
3310  
3315  
3320  
3325  
3330  
3335  
3340  
3345  
3350  
3355  
3360  
3365  
3370  
3375  
3380  
3385  
3390  
3395  
3400  
3405  
3410  
3415  
3420  
3425  
3430  
3435  
3440  
3445  
3450  
3455  
3460  
3465  
3470  
3475  
3480  
3485  
3490  
3495  
3500  
3505  
3510  
3515  
3520  
3525  
3530  
3535  
3540  
3545  
3550  
3555  
3560  
3565  
3570  
3575  
3580  
3585  
3590  
3595  
3600  
3605  
3610  
3615  
3620  
3625  
3630  
3635  
3640  
3645  
3650  
3655  
3660  
3665  
3670  
3675  
3680  
3685  
3690  
3695  
3700  
3705  
3710  
3715  
3720  
3725  
3730  
3735  
3740  
3745  
3750  
3755  
3760  
3765  
3770  
3775  
3780  
3785  
3790  
3795  
3800  
3805  
3810  
3815  
3820  
3825  
3830  
3835  
3840  
3845  
3850  
3855  
3860  
3865  
3870  
3875  
3880  
3885  
3890  
3895  
3900  
3905  
3910  
3915  
3920  
3925  
3930  
3935  
3940  
3945  
3950  
3955  
3960  
3965  
3970  
3975  
3980  
3985  
3990  
3995  
4000  
4005  
4010  
4015  
4020  
4025  
4030  
4035  
4040  
4045  
4050  
4055  
4060  
4065  
4070  
4075  
4080  
4085  
4090  
4095  
4100  
4105  
4110  
4115  
4120  
4125  
4130  
4135  
4140  
4145  
4150  
4155  
4160  
4165  
4170  
4175  
4180  
4185  
4190  
4195  
4200  
4205  
4210  
4215  
4220  
4225  
4230  
4235  
4240  
4245  
4250  
4255  
4260  
4265  
4270  
4275  
4280  
4285  
4290  
4295  
4300  
4305  
4310  
4315  
4320  
4325  
4330  
4335  
4340  
4345  
4350  
4355  
4360  
4365  
4370  
4375  
4380  
4385  
4390  
4395  
4400  
4405  
4410  
4415  
4420  
4425  
4430  
4435  
4440  
4445  
4450  
4455  
4460  
4465  
4470  
4475  
4480  
4485  
4490  
4495  
4500  
4505  
4510  
4515  
4520  
4525  
4530  
4535  
4540  
4545  
4550  
4555  
4560  
4565  
4570  
4575  
4580  
4585  
4590  
4595  
4600  
4605  
4610  
4615  
4620  
4625  
4630  
4635  
4640  
4645  
4650  
4655  
4660  
4665  
4670  
4675  
4680  
4685  
4690  
4695  
4700  
4705  
4710  
4715  
4720  
4725  
4730  
4735  
4740  
4745  
4750  
4755  
4760  
4765  
4770  
4775  
4780  
4785  
4790  
4795  
4800  
4805  
4810  
4815  
4820  
4825  
4830  
4835  
4840  
4845  
4850  
4855  
4860  
4865  
4870  
4875  
4880  
4885  
4890  
4895  
4900  
4905  
4910  
4915  
4920  
4925  
4930  
4935  
4940  
4945  
4950  
4955  
4960  
4965  
4970  
4975  
4980  
4985  
4990  
4995  
5000  
5005  
5010  
5015  
5020  
5025  
5030  
5035  
5040  
5045  
5050  
5055  
5060  
5065  
5070  
5075  
5080  
5085  
5090  
5095  
5100  
5105  
5110  
5115  
5120  
5125  
5130  
5135  
5140  
5145  
5150  
5155  
5160  
5165  
5170  
5175  
5180  
5185  
5190  
5195  
5200  
5205  
5210  
5215  
5220  
5225  
5230  
5235  
5240  
5245  
5250  
5255  
5260  
5265  
5270  
5275  
5280  
5285  
5290  
5295  
5300  
5305  
5310  
5315  
5320  
5325  
5330  
5335  
5340  
5345  
5350  
5355  
5360  
5365  
5370  
5375  
5380  
5385  
5390  
5395  
5400  
5405  
5410  
5415  
5420  
5425  
5430  
5435  
5440  
5445  
5450  
5455  
5460  
5465  
5470  
5475  
5480  
5485  
5490  
5495  
5500  
5505  
5510  
5515  
5520  
5525  
5530  
5535  
5540  
5545  
5550  
5555  
5560  
5565  
5570  
5575  
5580  
5585  
5590  
5595  
5600  
5605  
5610  
5615  
5620  
5625  
5630  
5635  
5640  
5645  
5650  
5655  
5660  
5665  
5670  
5675  
5680  
5685  
5690  
5695  
5700  
5705  
5710  
5715  
5720  
5725  
5730  
5735  
5740  
5745  
5750  
5755  
5760  
5765  
5770  
5775  
5780  
5785  
5790  
5795  
5800  
5805  
5810  
5815  
5820  
5825  
5830  
5835  
5840  
5845  
5850  
5855  
5860  
5865  
5870  
5875  
5880  
5885  
5890  
5895  
5900  
5905  
5910  
5915  
5920  
5925  
5930  
5935  
5940  
5945  
5950  
5955  
5960  
5965  
5970  
5975  
5980  
5985  
5990  
5995  
6000  
6005  
6010  
6015  
6020  
6025  
6030  
6035  
6040  
6045  
6050  
6055  
6060  
6065  
6070  
6075  
6080  
6085  
6090  
6095  
6100  
6105  
6110  
6115  
6120  
6125  
6130  
6135  
6140  
6145  
6150  
6155  
6160  
6165  
6170  
6175  
6180  
6185  
6190  
6195  
6200  
6205  
6210  
6215  
6220  
6225  
6230  
6235  
6240  
6245  
6250  
6255  
6260  
6265  
6270  
6275  
6280  
6285  
6290  
6295  
6300  
6305  
6310  
6315  
6320  
6325  
6330  
6335  
6340  
6345  
6350  
6355  
6360  
6365  
6370  
6375  
6380  
6385  
6390  
6395  
6400  
6405  
6410  
6415  
6420  
6425  
6430  
6435  
6440  
6445  
6450  
6455  
6460  
6465  
6470  
6475  
6480  
6485  
6490  
6495  
6500  
6505  
6510  
6515  
6520  
6525  
6530  
6535  
6540  
6545  
6550  
6555  
6560  
6565  
6570  
6575  
6580  
6585  
6590  
6595  
6600  
6605  
6610  
6615  
6620  
6625  
6630  
6635  
6640  
6645  
6650  
6655  
6660  
6665  
6670  
6675  
6680  
6685  
6690  
6695  
6700  
6705  
6710  
6715  
6720  
6725  
6730  
6735  
6740  
6745  
6750  
6755  
6760  
6765  
6770  
6775  
6780  
6785  
6790  
6795  
6800  
6805  
6810  
6815  
6820  
6825  
6830  
6835  
6840  
6845  
6850  
6855  
6860  
6865  
6870  
6875  
6880  
6885  
6890  
6895  
6900  
6905  
6910  
6915  
6920  
6925  
6930  
6935  
6940  
6945  
6950  
6955  
6960  
6965  
6970  
6975  
6980  
6985  
6990  
6995  
7000  
7005  
7010  
7015  
7020  
7025  
7030  
7035  
7040  
7045  
7050  
7055  
7060  
7065  
7070  
7075  
7080  
7085  
7090  
7095  
7100  
7105  
7110  
7115  
7120  
7125  
7130  
7135  
7140  
7145  
7150  
7155  
7160  
7165  
7170  
7175  
7180  
7185  
7190  
7195  
7200  
7205  
7210  
7215  
7220  
7225  
7230  
7235  
7240  
7245  
7250  
7255  
7260  
7265  
7270  
7275  
7280  
7285  
7290  
7295  
7300  
7305  
7310  
7315  
7320  
7325  
7330  
7335  
7340  
7345  
7350  
7355  
7360  
7365  
7370  
7375  
7380  
7385  
7390  
7395  
7400  
7405  
7410  
7415  
7420  
7425  
7430  
7435  
7440  
7445  
7450  
7455  
7460  
7465  
7470  
7475  
7480  
7485  
7490  
7495  
7500  
7505  
7510  
7515  
7520  
7525  
7530  
7535  
7540  
7545  
7550  
7555  
7560  
7565  
7570  
7575  
7580  
7585  
7590  
7595  
7600  
7605  
7610  
7615  
7620  
7625  
7630  
7635  
7640  
7645  
7650  
7655  
7660  
7665  
7670  
7675  
7680  
7685  
7690  
7695  
7700  
7705  
7710  
7715  
7720  
7725  
7730  
7735  
7740  
7745  
7750  
7755  
7760  
7765  
7770  
7775  
7780  
7785  
7790  
7795  
7800  
7805  
7810  
7815  
7820  
7825  
7830  
7835  
7840  
7845  
7850  
7855  
7860  
7865  
7870  
7875  
7880  
7885  
7890  
7895  
7900  
7905  
7910  
7915  
7920  
7925  
7930  
7935  
7940  
7945  
7950  
7955  
7960  
7965  
7970  
7975  
7980  
7985  
7990  
7995  
8000  
8005  
8010  
8015  
8020  
8025  
8030  
8035  
8040  
8045  
8050  
8055  
8060  
8065  
8070  
8075  
8080  
8085  
8090  
8095  
8100  
8105  
8110  
8115  
8120  
8125  
8130  
8135  
8140  
8145  
8150  
8155  
8160  
8165  
8170  
8175  
8180  
8185  
8190  
8195  
8200  
8205  
8210  
8215  
8220  
8225  
8230  
8235  
8240  
8245  
8250  
8255  
8260  
8265  
8270  
8275  
8280  
8285  
8290  
8295  
8300  
8305  
8310  
8315  
8320  
8325  
8330  
8335  
8340  
8345  
8350  
8355  
8360  
8365  
8370  
8375  
8380  
8385  
8390  
8395  
8400  
8405  
8410  
8415  
8420  
8425  
8430  
8435  
8440  
8445  
8450  
8455  
8460  
8465  
8470  
8475  
8480  
8485  
8490  
8495  
8500  
8505  
8510  
8515  
8520  
8525  
8530  
8535  
8540  
8545  
8550  
8555  
8560  
8565  
8570  
8575  
8580  
8585  
8590  
8595  
8600  
8605  
8610  
8615  
8620  
8625  
8630  
8635  
8640  
8645  
8650  
8655  
8660  
8665  
8670  
8675  
8680  
8685  
8690  
8695  
8700  
8705  
8710  
8715  
8720  
8725  
8730  
8735  
8740  
8745  
8750  
8755  
8760  
8765  
8770  
8775  
8780  
8785  
8790  
8795  
8800  
8805  
8810  
8815  
8820  
8825  
8830  
8835  
8840  
8845  
8850  
8855  
8860  
8865  
8870  
8875  
8880  
8885  
8890  
8895  
8900  
8905  
8910  
8915  
8920  
8925  
8930  
8935  
8940  
8945  
8950  
8955  
8960  
8965  
8970  
8975  
8980  
8985  
8990  
8995  
9000  
9005  
9010  
9015  
9020  
9025  
9030  
9035  
9040  
9045  
9050  
9055  
9060  
9065  
9070  
9075  
9080  
9085  
9090  
9095  
9100  
9105  
9110  
9115  
9120  
9125  
9130  
9135  
9140  
9145  
9150  
9155  
9160  
9165  
9170  
9175  
9180  
9185  
9190  
9195  
9200  
9205  
9210  
9215  
9220  
9225  
9230  
9235  
9240  
9245  
9250  
9255  
9260  
9265  
9270  
9275  
9280  
9285  
9290  
9295  
9300  
9305  
9310  
9315  
9320  
9325  
9330  
9335  
9340  
9345  
9350  
9355  
9360  
9365  
9370  
9375  
9380  
9385  
9390  
9395  
9400  
9405  
9410  
9415  
9420  
9425  
9430  
9435  
9440  
9445  
9450  
9455  
9460  
9465  
9470  
9475  
9480  
9485  
9490  
9495  
9500  
9505  
9510  
9515  
9520  
9525  
9530  
9535  
9540  
9545  
9550  
9555  
9560  
9565  
9570  
9575  
9580  
9585  
9590  
9595  
9600  
9605  
9610  
9615  
9620  
9625  
9630  
9635  
9640  
9645  
9650  
9655  
9660  
9665  
9670  
9675  
9680  
9685  
9690  
9695  
9700  
9705  
9710  
9715  
9720  
9725  
9730  
9735  
9740  
9745  
9750  
9755  
9760  
9765  
9770  
9775  
9780  
9785  
9790  
9795  
9800  
9805  
9810  
9815  
9820  
9825  
9830  
9835  
9840  
9845  
9850  
9855  
9860  
9865  
9870  
9875  
9880  
9885  
9890  
9895  
9900  
9905  
9910  
9915  
9920  
9925  
9930  
9935  
9940  
9945  
9950  
9955  
9960  
9965  
9970  
9975  
9980  
9985  
9990  
9995  
10000  
10005  
10010  
10015  
10020  
10025  
10030  
10035  
10040  
10045  
10050  
10055  
10060  
10065  
10070  
10075  
10080  
10085  
10090  
10095  
10100  
10105  
10110  
10115  
10120  
10125  
10130  
10135  
10140  
10145  
10150  
10155  
10160  
10165  
10170  
10175  
10180  
10185  
10190  
10195  
10200  
10205  
10210  
10215  
10220  
10225  
1023



antibodies will neutralize the growth promoting activity of KGF, thereby enabling mechanistic studies and, ultimately, therapy for clinical conditions involving excessive levels of KGF.

5

#### BRIEF DESCRIPTION OF THE DRAWINGS

Fig. I-1 depicts results of heparin-Sepharose affinity chromatography of conditioned medium from M426 human embryonic fibroblasts showing that greater than 90% of the mitogenic activity for mouse keratinocytes (BALB/MK) eluted with 0.6 M NaCl.

10 Fig. I-2 illustrates results of further purification of the mitogen from human fibroblasts using HPLC with an adsorptive matrix. Panel (A) shows the profile on reversed-phase ( $C_4$ ) HPLC of BALB/MK mitogenic activity. Panel (B) presents electrophoretic ( $\text{NaDodSO}_4$ /PAGE) analysis of selected fractions from the  $C_4$  chromatography shown in panel A, demonstrating that the peak HPLC fractions contained a single band on the silver stained gel. Panel (C) is a bar graph of DNA synthesis in BALB/MK cells triggered by the fractions analyzed in Panel B, showing that the relative mitogenic activity correlated well with the intensity of the protein band across the activity profile.

20 Fig. I-3 presents an alternative purification step to RP-HPLC, using sieving chromatography with a (TSK G3000SW GlasPac) column run in aqueous solution near physiologic pH, which resulted in a major peak of mitogenic activity in the BALB/MK bioassay.

25 Fig. I-4 illustrates a comparison of BALB/MK DNA synthesis in response to TSK-purified mitogen and other growth factors.

Fig. I-5 shows comparisons-of growth of BALB/MK cells in a chemically defined medium in response to different combinations of growth factors.

30 Table I-1 summarizes the results from various purification steps, documenting that sieving chromatography provided a far better recovery of activity than the adsorptive RP-HPLC approach.

Table I-2 recapitulates data on the target cell specificities of various growth factors, demonstrating that the newly isolated factor exhibited a strong mitogenic effect on keratinocytes (BALB/MK) and, in striking contrast, had no detectable effects on fibroblasts or human saphenous vein endothelial cells.

Fig. II-1 presents the nucleotide sequence and deduced amino acid sequence of KGF cDNA, as well as identification of RNAs transcribed from the KGF gene. Panel (A) outlines a schematic representation of human KGF cDNA clones. Panel (B) document the KGF cDNA nucleotide and predicted amino acid sequences. (C) Identification of RNA transcripts of KGF genes by Northern blot analysis.

Fig. II-2 illustrates the topological comparison of the FGF family of related molecules, including KGF, with emphasis on the two protein domains that sharp high homology, the putative signal peptide sequences, and the two conserved cysteine residues.

Fig. II-3 shows (Northern blot) analyses of expression of KGF-related mRNA in selected normal human cell lines and tissues, revealing that the predominant 2.4 kb transcript was present in RNA from stromal fibroblasts but was not detected in the epithelial cell lines.

Table II-1 summarizes a comparison of the effect of heparin on KGF mitogenic activity with effects on other growth factors, showing that thymidine incorporation into DNA by BALB/MK cells in response to KGF was inhibited by heparin, in contrast, to the activities of both aFGF and bFGF which were increased by the same treatment.

#### DESCRIPTION OF SPECIFIC EMBODIMENTS

This invention relates, in part, to purified KGF or KGF-like proteins and methods for preparing these proteins. A principal embodiment of this aspect of this invention relates to homogeneous KGF characterized by an apparent molecular weight of about 28 kDa based on migration in NaDodSO<sub>4</sub>/PAGE, movement as a single peak on

reversed-phase high performance liquid chromatography, and a specific activity of at least about  $3.4 \times 10^4$  units per milligram, and preferably at least about  $3.2 \times 10^5$  units per milligram, where one unit of activity is defined as that amount which causes half of the maximal possible stimulation of DNA synthesis in certain epithelial (keratinocyte) cells under standard assay conditions outlined below.

To identify novel growth factors specific for epithelial cell types, a clonal BALB/c mouse keratinocyte cell line, designated BALB/MK (Weissman, B.E. and Aaronson, S.A. (1983) *Cell* 32, 599-606) was employed as an indicator cell to detect such factors. These cells are dependent for their growth upon an exogenous source of an epithelial cell mitogen even in medium containing serum (Weissman, B.E. and Aaronson, S.A. (1983) *Cell* 32, 599-606). The development of chemically defined medium for these cells has made it possible to demonstrate that two major mitogenic pathways are required for BALB/MK proliferation. One involves insulin-like growth factor I (or insulin at high concentration) and the other is satisfied by epidermal growth factor (EGF), transforming growth factor  $\alpha$  (TGF $\alpha$ ), acidic fibroblast growth factor (aFGF) or basic fibroblast growth factor (bFGF) (Falco, J.P., Taylor, W.G., DiFiore, P.P., Weissman, B.E., and Aaronson, S.A. (1988) *Oncogene* 2, 573-578).

By using BALB/MK as the prototypical epithelial cell line and NIH/3T3 as its fibroblast counterpart, conditioned media from various human cell lines were assayed for new epithelial cell-specific mitogens. These bioassays of this invention enabled the purification to homogeneity of one such novel growth factor, released by a human embryonic lung fibroblast line, and designated herein as keratinocyte growth factor (KGF).

In brief, the bioassay for KGF-like activity under standard conditions comprises the following steps:

(i) Mouse keratinocytes (BALB/MK cells) are grown in culture to confluency and then maintained for 24-72 hr in serum-free medium;

(ii) Following addition of test samples, stimulation of DNA synthesis is determined by incorporation of <sup>3</sup>H-thymidine into acid-precipitable DNA.

To determine the target cell specificity of a mitogenic growth factor, the DNA synthesis stimulation, expressed as ratio of stimulated synthesis over background incorporation of thymidine in the absence of added test sample, can be compared to analogous stimulation observed in cells other than keratinocytes under the same assay conditions. In such comparisons, KGF mitogenic activity will exhibit marked specificity for the keratinocytes as opposed to fibroblasts (at least about 500-fold greater stimulation) and lesser but significant (at least about 50-fold) greater activity on keratinocytes than on other exemplary epithelial cell types (see Table I-2 for further data, and Materials and Methods in Experimental Section I for details of the standard conditions of the bioassay).

By employing a method of KGF production involving culturing cells and isolating mitogenic activity, which method comprises ultrafiltration, heparin-Sepharose affinity chromatography (HSAC) and adsorptive reversed-phase high performance liquid chromatography (RP-HPLC) or, alternatively, molecular sieving HPLC (TSK-HPLC), according to the present invention, a quantity was isolated sufficient to permit detailed characterization of the physical and biological properties of this molecule.

To summarize, the method for production of KGF from producing cells such as M426 human embryonic fibroblasts (Aaronson, S.A. and Todaro, G.J. (1968) *Virology* 36, 254-261), for example, comprises the following steps:

(i) Preparation of conditioned media (e.g., 10 liters) using monolayer cultures cycled from serum-containing to serum-free medium and storing the serum-free harvest at -70°C until further use;

(ii) Concentration by ultrafiltration using membranes having a 10 kDa molecular weight cutoff in several successive steps with intervening dilution in buffer (to

facilitate removal of low molecular weight materials), followed by optional storage at  $-70^{\circ}\text{C}$ ;

(iii) Affinity chromatography on heparin attached to a polymeric support (e.g., Sepharose) with elution by  
5 a gradient of increasing NaCl concentration;

(iv) Concentration by a factor of at least ten- to twenty-fold with small scale ultrafiltration devices with a 10 kDa molecular weight cutoff (e.g., a Centricon-10 microconcentrator from Amicon) and storage at  $-70^{\circ}\text{C}$ .

10 The next step of the purification process comprises either step (v) or, alternatively, step (vi), as follows:

(V) Reversed-phase HPLC of active fractions (0.6 M NaCl pool) from the previous HSAC step in organic solvent systems;

15 or,

(vi) Molecular sieve HPLC (e.g., on a TSK-G3000SW Glas-Pac Column from LKB) in aqueous buffer at near physiological pH (e.g., Tris-HCl, pH 6.8/0.5M NaCl) followed by storage at  $-70^{\circ}\text{C}$ .

20 A preparation made by the TSK step (vi) was almost as pure as one obtained from RP-HPLC, as judged by silver-stained NaDodSO<sub>4</sub>/PAGE (data not shown); but the TSK approach provided a far better recovery of activity (Table I-1). Further, the TSK-purified material had a  
25 higher specific activity than the RP-HPLC material. KGF prepared by the TSK procedure above stimulated DNA synthesis in epithelial cells at sub-nanomolar concentrations, but failed to induce any thymidine incorporation into DNA of fibroblasts or endothelial cells at  
30 comparable or higher concentrations (up to 5 nM). The activity was sensitive to acid, heat and solvents used in the RP-HPLC step. (See Experimental Section I for data on sensitivities and further details of the production method.)

35 Using standard methodology well known in the art, an unambiguous amino acid sequence was determined for positions 2-13 from the amino terminus of the purified KGF, as follows: Asn-Asp-Met-Thr-Pro-Glu-Gln-Met-Ala-Thr-Asn-Val (see Experimental Section I).

The present invention also includes DNA segments encoding KGF and KGF-like polypeptides. The DNAs of this invention are exemplified by DNAs referred to herein as: human cDNA clones 32 and 49 derived from polyadenylated  
5 RNA extracted from the human embryonic lung fibroblast cell line M426; recombinants and mutants of these clones; and related DNA segments which can be detected by hybridization to these DNA segments.

As described in Experimental Section II, to search  
10 for cDNA clones corresponding to the known portion of the KGF amino acid sequence, two pools of oligonucleotide probes were generated based upon all possible nucleotide sequences encoding the nine-amino acid sequence, Asn-Asp-Met-Thr-Pro-Glu-Gln-Met-Ala. A cDNA library was  
15 constructed in a cDNA cloning vector,  $\lambda$ pCEV9, using polyadenylated RNA extracted from the human embryonic lung fibroblast cell line M426 which was the initial source of the growth factor. Screening of the library ( $9 \times 10^5$  plaques) with the  $^{32}\text{P}$ -labelled oligonucleotides identified  
20 88 plaques which hybridized to both probes.

Of 10 plaque-purified clones that were analyzed, one, designated clone 49, had a cDNA insert of 3.5 kb, while the rest had inserts ranging from 1.8 kb to 2.1 kb. Analysis of the smaller clones revealed several common  
25 restriction sites, and sequencing of a representative smaller clone, designated clone 32, along with clone 49, demonstrated that they were overlapping cDNAs (Fig II-1A). Alignment of the two cDNAs established a continuous sequence of 3.85 kb containing the complete KGF coding  
30 sequence. The sense strand DNA nucleotide sequence, and the predicted primary protein sequence encoded, are shown for the full-length composite KGF cDNA sequence in Fig. II-1B.

These DNAs, cDNA clones 32 and 49, as well as  
35 recombinant forms of these segments comprising the complete KGF coding sequence, are most preferred DNAs of this invention.

From the cDNA sequence, it is apparent that the primary KGF, FGF-5 and *hst* translation products contain

hydrophobic N-terminal regions which likely serve as signal sequences, based on similarity to such sequences in a variety of other proteins. Accordingly, this N-terminal domain is not present in the purified mature KGF molecule which is secreted by human embryonic fibroblasts.

Furthermore, KGF shares with all other members of the FGF family two major regions of homology, spanning amino acids 65-156 and 162-189 in the predicted KGF sequence, which are separated by short, nonhomologous series of amino acids of various lengths in the different family members. The sequence of the purified form of KGF contains five cysteine residues, two of which are conserved throughout the family of FGF related proteins. Five pairs of basic residues occur throughout the KGF sequence. This same pattern has been observed in other FGF family members.

It should be obvious to one skilled in the art that, by using the DNAs and RNAs of this invention in hybridization methods (such as Southern blot analyses of genomic human DNAs), especially the most preferred DNAs listed herein above, without undue experimentation, it is possible to screen genomic or cDNA libraries to find other KGF-like DNAs which fall within the scope of this invention. Furthermore, by so using DNAs of this invention, genetic markers associated with the KGF gene, such as restriction fragment length polymorphisms (RFLPs), may be identified and associated with inherited clinical conditions involving this or other nearby genes.

This invention also includes modified forms of KGF DNAs. According to a chief embodiment of this aspect of the invention, such modified DNAs encode KGF-like proteins comprising segments of amino acid sequences of KGF and at least one other member of the FGF peptide family. Thus, for example, since there is no significant N-terminal homology between the secreted form of KGF and analogous positions in other FGF-related proteins, polypeptides with novel structural and functional properties may be created by grafting DNA segments encoding the

distinct N-terminal segments of another polypeptide in the FGF family onto a KGF DNA segment in place of its usual N-terminal sequence.

5 The polypeptide chimeras produced by such modified DNAs are useful for determining whether the KGF NH<sub>2</sub>-terminal domain is sufficient to account for its unique target cell specificity. Studies on chimeras should also provide insights into which domains contribute the different effects of heparin on their biologic  
10 activities.

Indeed, the utility of this approach has already been confirmed by the successful engineering and expression of a chimeric molecule in which 46 amino acids from the NH<sub>2</sub>-terminus of the secreted form of KGF (beginning with the  
15 amino terminal cys residue of the mature KGF form, numbered 32 in Fig II-1, and ending at KGF residue 78, arg) is linked to 116 amino acids of the C-terminal core of aFGF (beginning at residue 39, arg, and continuing to the C-terminal end of the aFGF coding sequence. This  
20 chimeric product has a target cell preference for keratinocytes, like KGF, but its activity is enhanced by heparin, a characteristic which parallels that of aFGF rather than KGF. This novel KGF-like growth factor may have advantages in clinical applications where  
25 administration of an epithelial-specific growth factor is desirable in the presence of heparin, a commonly used anticoagulant. Further details of the construction of this chimeric molecule and the properties of the polypeptide are described in Experimental Section II.

30 Other DNAs of this invention include the following recombinant DNA molecules comprising a KGF cDNA and any of the following exemplary vector DNAs: a bacteriophage  $\lambda$  cloning vector ( $\lambda$ pCEV9); a DNA sequencing plasmid vector (a pUC variant); a bacterial expression vector  
35 (pKK233-2); or a mammalian expression vector (pMMT/neo). Such recombinant DNAs are exemplified by constructs described in detail in the Experimental Sections.

Most preferred recombinant molecules include the following: molecules comprising the coding sequence for



the secreted form of KGF and a bacterial expression vector (e.g., pKK233-2) or a cDNA encoding the entire primary translation product (including the NH<sub>2</sub>-terminal signal peptide) and a mammalian expression vector (exemplified by pMMT) capable of expressing inserted DNAs in mammalian (e.g., NIH/3T3) cells.

Construction of recombinant DNAs containing KGF DNA and a bacterial expression vector is described in Experimental Section II. In brief, KGF cDNA was expressed to produce polypeptide in *E. coli* by placing its coding sequence under control of the hybrid trk promoter in the plasmid expression vector pKK233-2 (Amman, E. and Brosius, J. (1985) *Gene* 40, 183).

Construction of recombinant DNAs comprising KGF DNA and a mammalian vector capable of expressing inserted DNAs in cultured human or animal cells, can be carried out by standard gene expression technology using methods well known in the art for expression of such a relatively simple polypeptide. One specific embodiment of a recombinant DNA of this aspect of the present invention, involving the mammalian vector pMMT, is described further below in this section under recombinant cells of this invention.

DNAs and sense strand RNAs of this invention can be employed, in conjunction with protein production methods of this invention, to make large quantities of substantially pure KGF or KGF-like proteins. Substantially pure KGF protein thus produced can be employed, using well-known techniques, in diagnostic assays to determine the presence of receptors for this protein in various body fluids and tissue samples.

Accordingly, this invention also comprises a cell, preferably a bacterial or mammalian cell, transformed with a DNA of the invention, wherein the transforming DNA is capable of being expressed. In a preferred embodiment of this aspect of the invention, the cell transformed by the DNA of the invention produces KGF protein in a fully mitogenic form. Most preferably, these proteins will be of a secreted form (i.e., lacking an apparent signal

sequence). These protein factors can be used for functional studies, and can be purified for additional biochemical and functional analyses, such as qualitative and quantitative receptor binding assays.

5        Recombinant *E. coli* cells have been constructed in a bacterial expression vector, pKK233-2, for production of KGF, as detailed in Experimental Section II. In summary, several recombinant bacterial clones were tested for protein production by the usual small scale methods.  
10       All recombinants tested synthesized a protein that was recognized by antibodies raised against an amino-terminal KGF peptide (see below). One recombinant was grown up in a one liter culture which produced recombinant KGF that efficiently stimulated thymidine incorporation into DNA  
15       of BALB/MK keratinocyte cells, but was only marginally active on NIH/3T3 fibroblasts. Half-maximal stimulation of the BALB/MK cells in the standard keratinocyte bioassay was achieved with a concentration of between 2 to 5 ng/ml, compared to a concentration of 10 to 15 ng/ml  
20       for KGF purified from M426 cells.

      One liter of bacterial cells yielded approximately 50  $\mu$ g of Mono-S purified recombinant KGF. It will be apparent to those skilled in the art of gene expression that this initial yield can be improved substantially  
25       without undue experimentation by application of a variety known recombinant DNA technologies.

      Recombinant mammalian (NIH/3T3 mouse) cells have also been constructed using the entire KGF cDNA coding sequence (including the NH<sub>2</sub>-terminal signal peptide) and  
30       the vector pMMT/neo, which carries mouse metallothioneine (MMT) promoter and the selective marker gene for neomycin resistance. The cells are being evaluated for KGF production, particularly for secretion of the mature form (lacking signal peptide) produced by human fibroblasts,  
35       using bioassays of the present invention. This same vector and host cell combination has been used successfully to express several other similar recombinant polypeptides, including high levels of Platelet-Derived Growth Factor (PDGF) A and B chains (Sakai, R.K., Scharf,

S., Faloona, F., Mullis, K.B., Norn, G.T., Erlich, H.A. and Arnheim, N. (1985) *Science* 230, 1350-1354). Accordingly, it will be recognized by those skilled in the art that high yields of recombinant KGF can be achieved in this manner, using the aforementioned recombinant DNAs and transformed cells of this invention.

Ultimately, large-scale production can be used to enable clinical testing in conditions requiring specific stimulation of epithelial cell growth. Materials and methods for preparing pharmaceutical compositions for administration of polypeptides topically (to skin or to the cornea of the eye, for example) or systemically are well known in the art and can be adapted readily for administration of KGF and KGF-like peptides without undue experimentation.

This invention also comprises novel antibodies made against a peptide encoded by a DNA segment of the invention. This embodiment of the invention is exemplified by several kinds of antibodies which recognize KGF. These have been prepared using standard methodologies well known in the art of experimental immunology, as outlined in Experimental Section II. These antibodies include: monoclonal antibodies raised in mice against intact, purified protein from human fibroblasts; polyclonal antibodies raised in rabbits against synthetic peptides with sequences based on amino acid sequences predicted from the KGF cDNA sequence [exemplified by a peptide with the sequence of KGF residues 32-45 plus an R at its C-terminus, namely, NDMTPEQMATNVR (using standard one-letter code for amino acid sequences; see Fig. II-1)]; polyclonal antibodies raised in rabbits against both naturally secreted KGF from human fibroblasts and recombinant KGF produced in *E. coli* (see above).

All tested antibodies recognize the recombinant as well as the naturally occurring KGF, either in a solid-phase (ELISA) assay and/or in a Western blot. Some exemplary antibodies, which are preferred antibodies of this invention, appear to neutralize mitogenic activity of KGF in the BALB/MK bioassay.

Fragments of antibodies of this invention, such as Fab or F(ab)' fragments, which retain antigen binding activity and can be prepared by methods well known in the art, also fall within the scope of the present invention. Further, this invention comprises pharmaceutical compositions of the antibodies of this invention, or active fragments thereof, which can be prepared using materials and methods for preparing pharmaceutical compositions for administration of polypeptides that are well known in the art and can be adapted readily for administration of KGF and KGF-like peptides without undue experimentation.

These antibodies, and active fragments thereof, can be used, for example, for detection of KGF in bioassays or for purification of the protein factors. They may also be used in approaches well known in the art, for isolation of the receptor for KGF, which, as described in Experimental Section II, appears to be distinct from those of all other known growth factors.

Those preferred antibodies, and fragments and pharmaceutical compositions thereof, which neutralize mitogenic activity of KGF for epithelial cells, as indicated by the BALB/MK assay, for instance, may be used in the treatment of clinical conditions characterized by excessive epithelial cell growth, including dysplasia and neoplasia (e.g., psoriasis, or malignant or benign epithelial tumors).

This invention further comprises novel bioassay methods for detecting the expression of genes related to DNAs of the invention. In some exemplary embodiments, DNAs of this invention were used as probes to determine steady state levels of related mRNAs. Methods for these bioassays of the invention, using KGF DNAs, and standard Northern blotting techniques, are described in detail in Experimental Section II.

One skilled in the art will recognize that, without undue experimentation, such methods may be readily applied to analysis of gene expression for KGF-like proteins, either in isolated cells or various tissues. Such bioassays may be useful, for example, for

identification of various classes of tumor cells or genetic defects in the epithelial growth processes.

Without further elaboration, it is believed that one of ordinary skill in the art, using the preceding description, and following the methods of the Experimental Sections below, can utilize the present invention to its fullest extent. The material disclosed in the Experimental Sections, unless otherwise indicated, is disclosed for illustrative purposes and therefore should not be construed as being limitive in any way of the appended claims.

#### EXPERIMENTAL SECTION I

##### IDENTIFICATION AND CHARACTERIZATION OF A NOVEL GROWTH FACTOR SPECIFIC FOR EPITHELIAL CELLS

This section describes experimental work leading to identification of a growth factor specific for epithelial cells in conditioned medium of a human embryonic lung fibroblast cell line. The factor, provisionally termed keratinocyte growth factor (KGF) because of its predominant activity on this cell type, was purified to homogeneity by a combination of ultrafiltration, heparin-Sepharose affinity chromatography and hydrophobic chromatography on a C<sub>4</sub> reversed-phase HPLC column, according to methods of this invention. KGF was found to be both acid and heat labile, and consisted of a single polypeptide chain with an apparent molecular weight of approximately 28,000 daltons. Purified KGF was a potent mitogen for epithelial cells, capable of stimulating DNA synthesis in quiescent BALB/MK epidermal keratinocytes by more than 500-fold with activity detectable at 0.1 nM and maximal at 1.0 nM. Lack of mitogenic activity on either fibroblasts or endothelial cells indicated that KGF possessed a target cell specificity distinct from any previously characterized growth factor. Microsequencing revealed an amino-terminal sequence containing no significant homology to any known protein. The release of this novel growth factor by human embryonic fibro-

blasts indicates that KGF plays a role in mesenchymal stimulation of normal epithelial cell proliferation.

#### METHODS AND MATERIALS

5        Preparation of Conditioned Media. An early passage  
of M426 human embryonic fibroblasts (Aaronson, S.A. and  
Todaro, G.J. (1968) *Virology* 36, 254-261) was plated onto  
175 cm<sup>2</sup> T-flasks and grown to confluence over 10-14 days  
in Dulbeccos modified Eagle's medium (DMEM; GIBCO)  
supplemented with 10% calf serum (GIBCO). Once  
10        confluent, the monolayers were cycled weekly from serum-  
containing to serum-free medium, the latter consisting of  
DMEM alone. The cells were washed twice with 5 ml of  
phosphate buffered saline prior to addition of 20 ml of  
DMEM. After 72 hrs, culture fluids were collected and  
15        replaced with 35 ml of serum-containing medium. The  
conditioned medium was stored at -70°C until further use.

20        Ultrafiltration. Approximately ten liters of  
conditioned medium were thawed, prefiltered through a  
0.50 micron filter (Millipore HAWP 142 50) and concen-  
trated to 200 ml using the Pellicon cassette system  
(Millipore XX42 00K 60) and a cassette having a 10 kDa  
molecular weight cutoff (Millipore PTGC 000 05). After  
concentration, the sample was subjected to two successive  
rounds of dilution with one liter of 20 mM Tris-HCl, pH  
25        7.5/0.3M NaCl, each followed by another step of ultra-  
filtration with the Pellicon system. Activity recovered  
in the retentate was either immediately applied to  
heparin-Sepharose resin or stored at -70°C.

30        Heparin-Sepharose Affinity Chromatography (HSAC).  
The retentate from ultrafiltration was loaded onto  
heparin-Sepharose resin (Pharmacia) which had been  
equilibrated in 20 mM Tris-HCl, pH 7.5/0.3 M NaCl. The  
resin was washed extensively until the optical density  
had returned to baseline and then subjected to a linear-  
35        step gradient of increasing NaCl concentration. After  
removing aliquots from the fractions for the thymidine  
incorporation bioassay, selected fractions were

concentrated ten- to twenty-fold with a Centricon-10 microconcentrator (Amicon) and stored at -70°C.

5        Reversed-Phase HPLC (RP-HPLC). Active fractions (0.6 M NaCl pool) from the HSAC were thawed, pooled and further concentrated with the Centricon-10 to a final  
10        volume of  $\leq 200 \mu\text{l}$ . The sample was loaded onto a Vydac C<sub>18</sub> HPLC column (The Separations Group, Hesperia, CA) which had been equilibrated in 0.1% trifluoroacetic acid (TFA, Fluka)/20% acetonitrile (Baker, HPLC grade) and eluted  
15        with a linear gradient of increasing acetonitrile. Aliquots for the bioassay were immediately diluted in a 10-fold excess of 50  $\mu\text{g/ml}$  BSA (Fraction V, Sigma)/20 mM Tris-HCl, pH 7.5. The remainder of the sample was dried in a Speed-Vac (Savant) in preparation for structural analysis.

Molecular Sieve HPLC. Approximately 50  $\mu\text{l}$  of the twice concentrated heparin-Sepharose fractions were loaded onto a TSK-G3000SW Glas-Pac Column (LKB) which had been equilibrated in 20 mM Tris-HCl, pH 6.8/0.5M NaCl.  
20        The sample was eluted in this buffer at a flow rate of 0.4 ml/min. After removing aliquots for the bioassay, the fractions were stored at -70°C.

NaDodSO<sub>4</sub>-Polyacrylamide Gel Electrophoresis (NaDodSO<sub>4</sub>/PAGE). Polyacrylamide gels were prepared with  
25        NaDodSO<sub>4</sub> according to the procedure of Laemmli (Laemmli, U.K. (1970) Nature 227, 680-685). Samples were boiled for 3 min in the presence of 2.5% 2-mercaptoethanol (vol/vol). The gels were fixed and stained with silver (Merril, C.R., Goldman, D., Sedman, S.A. and Ebert, M.H.  
30        (1981) Science 211, 1437-1438) using the reagents and protocol from BioRad. Molecular weight markers were from Pharmacia.

DNA Synthesis Stimulation. Ninety-six well micro-liter plates (Falcon No. 3596) were precoated with human  
35        fibronectin (Collaborative Research) at 1  $\mu\text{g/cm}^2$  prior to seeding with BALB/MK cells. Once confluent, the cells were maintained for 24-72 hr in serum-free medium containing 5  $\mu\text{g/ml}$  transferrin (Collaborative Research) and 30 nM Na<sub>2</sub>SeO<sub>3</sub> (Baker). Incorporation of <sup>3</sup>H-thymidine

(5  $\mu$ m/ml final concentration, NEN) into DNA was measured during a 6 hr period beginning at 16 hrs following addition of samples. The assay was terminated by washing the cells once with ice cold phosphate-buffered saline and twice with 5% trichloroacetic acid. The precipitate  
5 was redissolved in 0.25 M NaOH, transferred into liquid scintillation fluid (Biofluor, NEN) and counted.

Stimulation of DNA synthesis was monitored as described above for BALB/MK cells on a variety of other  
10 cell lines. NIH/3T3 fibroblasts (Jainchill, J.L., Aaronson, S.A. and Todaro, G.J. (1969) *J. Virol.* 4, 549-553) were available from the National Institutes of Health, while CCL208 Rhesus monkey bronchial epithelial cells (Caputo, J.L., Hay, R.J. and Williams, C.D. (1979)  
15 *In Vitro* 15, 222-223) were obtained from the American Type Culture Collection. The B5/589 human mammary epithelial cell line, prepared as described in (Stampfer, M.R. and Bartley, J.C. (1985) *Proc. Natl. Acad. Sci. USA* 82, 2394-2398), was obtained from Martha Stampfer  
20 (Lawrence Berkeley Laboratory). The mammary cells were grown in RPMI 1640 supplemented with 10% fetal calf serum and 4 ng/ml EGF. When maintained in serum-free conditions, the basal medium was DMEM. Primary cultures of human saphenous vein endothelial cells were prepared and  
25 maintained as described elsewhere (Sharefkin, J.B., Fairchild, K.D., Albus, R.A., Cruess, D.F. and Rich, N.M. (1986) *J. Surgical Res.* 41, 463-472). Epidermal growth factor and insulin were from Collaborative Research. Acidic FGF and bFGF were obtained from California  
30 Biotechnology, Inc. Recombinant TGF $\alpha$  was obtained from Genentech, Inc. Media and serum were either from GIBCO, Biofluids, Inc. or the NIH media unit.

Proliferation Assay. Thirty-five mm culture dishes were precoated sequentially with poly-D-lysine  
35 (20  $\mu$ g/cm<sup>2</sup>) (Sigma) and human fibronectin, and then seeded with approximately  $2.5 \times 10^4$  BALB/MK cells. The basic medium was a 1:1 mixture of Eagle's low Ca 2+ minimal essential medium and Ham's F-12 medium, supplemented with 5  $\mu$ g/ml transferrin, 30 nM Na<sub>2</sub>SeO<sub>3</sub>, and



0.2 mM ethanolamine (Sigma). Medium was changed every 2 or 3 days. After 10 days, the cells were fixed in formalin (Fisher Scientific Co.) and stained with Giemsa (Fisher Scientific Co.).

5        Protein microsequencing.        Approximately 4  $\mu$ g  
(~150 pmol) of protein from the active fractions of the  $C_4$  column were redissolved in 50% TFA and loaded onto an Applied Biosystems gas-phase protein sequenator. Twenty  
10        rounds of Edman degradation were carried out and identifications of amino acid derivatives were made with an automated on-line HPLC (Model 120A, Applied Biosystems).

### RESULTS

Growth Factor Detection and Isolation. Preliminary  
15        screening of conditioned media from various cell lines indicated that media from some fibroblast lines contained mitogenic activities detectable on both BALB/MK and NIH/3T3 cells. Whereas boiling destroyed the activity on BALB/MK, mitogenic activity on NIH/3T3 remained intact. Based on the known heat stability of EGF (Cohen, S.  
20        (1962) *J. Biol. Chem.* 237, 1555-1562) and TGF $\alpha$  (DeLarco, J.E. and Todaro, G.J. (1978) *Proc. Natl. Acad. Sci. USA* 75, 4001-4005), it was reasoned that the BALB/MK mitogenic activity might be due to an agent different from these known epithelial growth factors.

25        M426, a human embryonic lung fibroblast line, was selected as the most productive source of this activity for purification of the putative growth factor(s). Ultrafiltration with the Pellicon system provided a  
30        convenient way of reducing the sample volume to a suitable level for subsequent chromatography. Various combinations of sieving, ion exchange and isoelectric focusing chromatography were tried during the development of a purification scheme, but all resulted in unacceptably low yields. On the other hand, heparin-  
35        Sepharose affinity chromatography (HSAC), which has been employed in the purification of other growth factors (Raines, E. W. and Ross, R. (1982) *J. Biol. Chem.* 257, 5154-5160; Shing, Y., Folkman, J., Sullivan, R.,

Butterfield, C., Murray, J. and Klagsburn, M. (1984) *Science* 223, 1296-1299; Gospodarowicz, D., Cheng, J., Lui, G.-M., Baird, A. and Bohlen, P. (1984) *Proc. Natl. Acad. Sci. USA* 81, 6963-6967; Maciag, T., Mehlman, T.,  
5 Friesel, R. and Schreiber, A.B. (1984) *Science* 225, 932-935; Conn, G. and Hatcher, V.B. (1984) *Biochem. Biophys. Res. Comm.* 124, 262-268; Lobb, R.R. and Fett, J.W. (1984) *Biochemistry* 23, 6295-6299, proved to be useful as an  
10 early purification step in the present invention. While estimates of recovered specific activity were uncertain at this stage because of the likely presence of other factors, the apparent yield-of activity was 50-70% with a corresponding enrichment of approximately 1000 fold.

As shown in Fig. I-1, greater than 90% of the BALB/MK  
15 mitogenic activity eluted from the HSAC column with 0.6M NaCl. This peak of activity was not associated with any activity on NIH/3T3 cells (data not shown). A much smaller peak of BALB/MK mitogenic activity consistently emerged with 0.8 - 1.2M NaCl.

20 Due to the reproducibility of the HSAC pattern, active fractions could be identified presumptively on the basis of the gradient and optical density profile. Prompt concentration of 10-20 fold with the Centricon-10 was found to be essential for stability, which could be  
25 maintained subsequently at -70°C for several months.

Final purification was achieved by RP-HPLC with a C<sub>4</sub> Vydac column, a preparative method suitable for amino acid sequence analysis. While the yield of activity from the C<sub>4</sub> step was usually only a few percent, this loss  
30 could be attributed to the solvents employed. In other experiments, exposure to 0.1% TFA/50% acetonitrile for 1 hr at room temperature reduced the mitogenic activity of the preparation by 98%. Nonetheless, as shown in Fig. I-2, a single peak of BALB/MK stimulatory activity  
35 was obtained, coinciding with a distinct peak in the optical density profile. The peak fractions produced a single band upon NaDodSO<sub>4</sub>/PAGE and silver staining of the gel (Fig. I-2B), and the relative mitogenic activity of

each tested fraction (Fig. I-2C) correlated well with the intensity of the bands across the activity profile.

5 An alternative purification step to the HPLC technique described above, using sieving chromatography with a TSK G3000SW GlasPac column run in aqueous solution near physiologic pH, resulted in a major peak of activity in the BALB/MK bioassay (Fig. I-3). This preparation was almost as pure as the one obtained from RP-HPLC as judged by silver-stained NaDodSO<sub>4</sub>/PAGE (data not shown) but  
10 provided a far better recovery of activity (Table I-1). The TSK-purified material was used routinely for biological studies as it had a higher specific activity.

In both types of purified preparations (i.e., purified by HPLC or molecular sieving), the profile of  
15 mitogenic activity was associated with a distinct band on NaDodSO<sub>4</sub>/PAGE which appeared to be indistinguishable in the two preparations.

Physical and Biological Characterization of the Growth Factor. The purified factor had an estimated  
20 molecular weight of about 28 kDa based on NaDodSO<sub>4</sub>/PAGE under reducing (Fig. I-2) and non-reducing conditions (data not shown). This value was in good agreement with its elution position on two different sizing columns run in solvents expected to maintain native conformation  
25 (TSK-G3000-SW, Fig. I-3, and superose-12, data not shown). From these data, the mitogen appears to consist of a single polypeptide chain with a molecular weight of 25-30 kDa.

The heat and acid lability of the mitogenic activity  
30 were demonstrated using the BALB/MK mitogenesis bioassay. While activity was unaffected by a 10 min incubation at 50°C, it was reduced by 68% after 10 min at 60°C and was undetectable after 3 min at 100°C. Exposure to 0.5M acetic acid for 60 min at room temperature resulted in a  
35 decline in activity to 14% of the control. In comparison, the mitogenic activity of the known growth factor, EGF, was not diminished by any of these treatments.

The dose response curve for the purified growth factor depicted in Fig. I-4 illustrates that as little as

0.1 nM led to a detectable stimulation of DNA synthesis. Thus, the activity range was comparable to that of the other growth factors analyzed to date. A linear relationship was observed in the concentration range  
5 0.1 - 1.0 nM with maximal stimulation of 600 fold observed at 1.0 nM. The novel factor consistently induced a higher level of maximal thymidine incorporation than EGF, aFGF, or bFGF in the BALB/MK keratinocytes (Fig. I-4).

10 The distinctive target cell specificity of this factor was demonstrated by comparing its activities on a variety of cell types with those of other growth factors known to possess epithelial cell mitogenic activity. As shown in Table I-2, the newly isolated factor exhibited  
15 a strong mitogenic effect on BALB/MK but also induced demonstrable incorporation of thymidine into DNA of the other epithelial cells tested. In striking contrast, the factor had no detectable mitogenic effects on mouse (or human, data not shown) fibroblasts or human saphenous  
20 vein endothelial cells.

By comparison, none of the other known growth factors appeared to preferentially stimulate keratinocytes. TGF $\alpha$  and EGF showed potent activity on fibroblasts, while the FGFs were mitogenic for endothelial cells as well as  
25 fibroblasts (Table I-2). Because of its specificity of epithelial cells and the sensitivity of keratinocytes in particular, the novel mitogen was provisionally designated as keratinocyte growth factor (KGF).

To establish that KGF not only would stimulate DNA  
30 synthesis but would also support sustained cell growth, the ability of BALB/MK cells to grow in a fully-defined, serum-free medium supplemented with this growth factor was assessed. As shown in Fig. I-5, KGF served as an excellent substitute for EGF but not insulin (or insulin-  
35 like growth factor I) in this chemically defined medium. Thus, KGF appears to act through the major signalling pathway shared by EGF, aFGF and bFGF for proliferation of BALB/MK cells.

Microsequencing Reveals a Unique N-terminal Amino Acid Sequence of KGF. To further characterize the growth factor, approximately 150 pmol of C<sub>4</sub>-purified material were subjected to amino acid sequence analysis. A single sequence was detected with unambiguous assignments made for cycles 2-13, as follows: X-Asn-Asp-Met-Thr-Pro-Glu-Gln-Met-Ala-Thr-Asn-Val. High background noise precluded an assignment for the first position which is, therefore, indicated by an X.

A computer search using the FASTP program (Lipman, D.J. and Pearson, R.W. (1985) *Science* 227, 1435-1441) revealed that the N-terminal amino acid sequence of KGF showed no significant homology to any protein in the National Biomedical Research Foundation data bank, thus supporting the novelty of this epithelial growth factor.

#### DISCUSSION

The studies described in this Experimental Section identified a human growth factor which has a unique specificity for epithelial cells. By employing ultra-filtration, HSAC and RP-HPLC or TSK sieving chromatography according to the present invention, a quantity sufficient to permit detailed characterization of the physical and biological properties of this molecule was isolated.

A single silver-stained band corresponding to a molecular weight of about 28,000 daltons was detected in the active fractions from RP-HPLC, and the intensity of the band was proportional to the level of mitogenic activity in these fractions. A band indistinguishable from that obtained by RP-HPLC was seen in the active fractions from TSK chromatography. The purified protein stimulated DNA synthesis in epithelial cells at sub-nanomolar concentrations, but failed to induce any thymidine incorporation in fibroblasts or endothelial cells at comparable or higher concentrations (up to 5 nM). This distinctive target cell specificity combined with the single novel N-terminal amino acid sequence

determined from the purified molecule lead to the conclusion that KGF represents a new growth factor.

In a chemically defined medium the purified factor was able to complement the insulin-like growth factor I/insulin growth requirement of BALB/MK cells and therefore must act through a signal transduction pathway shared with EGF, TGF $\alpha$  and the FGFs. Moreover, the new factor was more potent than any of the known epithelial cell mitogens in stimulating thymidine incorporation in BALB/MK cells. Preliminary evidence indicates that this factor is also capable of supporting proliferation of secondary cultures of human keratinocytes (data not shown).

Handling and storage of KGF were problematical during its purification. Besides its inherent lability to acid and heat, it was unstable to lyophilization or dialysis. After HSAC, complete loss of activity occurred within 24 hr despite the use of carrier proteins, heparin, protease inhibitors, siliconized tubes or storage at either 4° or -20°C. Only concentrating the sample at this stage could preserve its activity.

Furthermore, in order to transfer the dried, purified factor it was necessary to utilize either strong acid or detergent, consistent with an adsorptive tendency or insolubility. Thus, for preservation of activity, the purified factor was maintained in solution at high concentrations at -70°C where it remained stable for several months.

The ability of KGF to bind heparin may signify a fundamental property of this factor that has a bearing on its function *in vivo*. Growth factors with heparin-binding properties include aFGF (Maciag, T., Mehlman, T., Friesel, R. and Schreiber, A.B. (1984) *Science* 225, 932-935; Conn, G. and Hatcher, V.B. (1984) *Biochem. Biophys. Res. Comm.* 124, 262-268; Lobb, R.R. and Fett, J.W. (1984) *Biochemistry* 23, 6295-6299), bFGF (Gospodarowicz, D., Cheng, J., Lui, G.-M., Baird, A. and Bohlen, P. (1984) *Proc. Natl. Acad. Sci. USA* 81, 6963-6967, Lobb, R.R. and Fett, J.W. (1984) *Biochemistry* 23, 6295-6299) granulo-

cyte/macrophage colony stimulating factor (Roberts, R.,  
Gallagher, J., Spooncer, E., Allen, T.D., Bloomfield, F.  
and Dexter, T. M. (1988) *Nature* 332, 376-378) and inter-  
leukin 3 (Roberts, R., Gallagher, J., Spooncer, E.,  
5 Allen, T.D., Bloomfield, F. and Dexter, T. M. (1988)  
*Nature* 332, 376-378). Each of these is produced by  
stromal cells (Roberts, R., Gallagher, J., Spooncer, E.,  
Allen, T.D., Bloomfield, F. and Dexter, T. M. (1988)  
*Nature* 332, 376-378; Libermann, T.A., Friesel, R., Jaye,  
10 M., Lyall, R. M., Westermarck, B., Drohen, W., Schmidt,  
A., Maciag, T. and Schlessinger, J. (1987) *EMBOJ.*, 61  
1627-1632; Shipley, G. D., Sternfeld, M. D., Coffey, R.  
J. and Pittelkow, M. R. (1988) *J. Cell Biochem. Supp* 12A,  
125, abstr. C420). Such factors appear to be deposited  
15 in the extracellular matrix, or on proteoglycans coating  
the stromal cell surface (Roberts, R., Gallagher, J.,  
Spooncer, E., Allen, T.D., Bloomfield, F. and Dexter, T.  
M. (1988) *Nature* 332, 376-378, Vlodavsky, I., Folkman,  
J., Sullivan, R., Fridman, R., Ishai-Michaeli, R., Sasse,  
20 J. and Klagsburn, M. (1987) *Proc. Natl. Acad. Sci. USA*  
84, 2292-2296). It has been postulated that their  
storage, release and contact with specific target cells  
are regulated by this interaction (Roberts, R.,  
Gallagher, J., Spooncer, E., Allen, T.D., Bloomfield, F.  
25 and Dexter, T. M. (1988) *Nature* 332, 376-378, Vlodavsky,  
I., Folkman, J., Sullivan, R., Fridman, R., Ishai-  
Michaeli, R., Sasse, J. and Klagsburn, M. (1987) *Proc.*  
*Natl. Acad. Sci. USA* 84, 2292-2296). While mesenchymal-  
derived effectors of epithelial cell proliferation have  
30 also been described (Gilchrest, B.A., Karassik, R.L.,  
Wilkins, L.M., Vrabel, M.A. and Maciag, T. (1983) *J. Cell*  
*Physiol.* 117, 2325-240, Chan, K. Y. and Haschke, R. H.  
(1983) *Exp. Eye Res.* 36, 231-246, Stiles, A. D., Smith,  
B. T. and Post, M. (1986) *Exp. Lung Res.* 11, 165-177),  
35 their identities have not been elucidated. Its heparin-  
binding properties, release by human embryonic fibroblast  
stromal cells, and epithelial cell tropism provide KGF  
with all of the properties expected of such a paracrine  
mediator of normal epithelial cell growth.

The partial amino acid sequence determined for this new growth factor has enabled molecular cloning of its coding sequence and determination of its structural relationship to known families of growth factors, as described in Experimental Section II, below.

## EXPERIMENTAL SECTION II

### CDNA SEQUENCE OF A NOVEL EPITHELIAL CELL SPECIFIC GROWTH FACTOR DEFINES A NEW MEMBER OF THE FGF FAMILY

10        Work in the previous Experimental Section I identified and purified a novel heparin-binding growth factor, designated keratinocyte growth factor (KGF), which is particularly active on keratinocytes and appears to be specific for epithelial cells. This second  
15        Experimental Section describes the isolation and characterization of cDNA clones encoding KGF, using synthetic oligonucleotides, based upon the experimentally determined NH<sub>2</sub>-terminal amino acid sequence, as hybridization probes. Nucleotide sequence analysis  
20        identified a 582-bp open reading frame which would code for a 194-amino acid polypeptide that is between 41% and 33% identical to the heparin-binding acidic and basic fibroblast growth factors (FGFs), and the related products of the *hst*, FGF-5 and *int-2* oncogenes. The KGF  
25        gene RNA transcript is expressed in normal fibroblasts of both embryonic and adult origin, but not in epithelial, endothelial or glial cells. Thus, KGF appears to be normally expressed by the mesenchyme, indicating a role in the regulation of epithelial cell proliferation.



## MATERIALS AND METHODS

Isolation of cDNA clones. The purification and N-terminal sequencing of KGF has been previously described (see Experimental Section I, above and Rubin, J.S., Osada, H., Finch, P.W., Taylor, W.G., Rudikoff, S. and Aaronson, S.A. (1989) *Proc. Natl. Acad. Sci. USA* (in press), February, 1989). Pools (50 pmole) of deoxyoligonucleotides described under Results were 5' end-labelled using 83 pmole of  $\gamma$ -<sup>32</sup>P-ATP (3000 Ci/mmole, Amersham) and 10 units of T4 polynucleotide kinase. The recombinant phage carrying cDNA clones were replica plated onto nitrocellulose filters and hybridized with <sup>32</sup>P-labelled deoxyoligonucleotides in 20% formamide, 10% dextran sulphate, 10 mM Tris-HCl (pH 7.5), 8 x SSC, 5x Denhardt's and 50 µg/ml denatured salmon sperm DNA, overnight at 42°C. Filters were washed in 0.5 x SSC, 0.1% SDS at 50°C and exposed to Kodak X-omat AR film.

DNA sequencing. The nucleotide sequence of the KGF cDNA was determined by the dideoxy chain termination method (Sanger, F., Nicklen, S. and Coulson, A.R. (1977) *Proc. Natl. Acad. Sci. USA* 74, 5463-5467), of overlapping restriction fragments, subcloned into pUC vectors (Yanisch-Perron, C., Vieira, J. and Messing, J. (1985) *Gene* 33, 103-119)

Construction of a bacterial expression vector for KGF cDNA. KGF cDNA encoding the mature, secreted form of the polypeptide was placed under control of the hybrid *trk* promoter in the plasmid expression vector pKK233-2 (Amman, E. and Brosius, J. (1985) *Gene* 40, 183), as follows. To accomplish this, a specific length of KGF cDNA that contained the information to code for the mature KGF molecule (i.e., without its signal peptide) was amplified using the polymerase chain reaction (PCR) technique (Sakai, R.K., Scharf, S., Faloona, F., Mullis, K.B., Norn, G.T., Erlich, H.A. and Arnheim, N. (1985) *Science* 230, 1350-1354). The fragment was directionally inserted between two sites in the vector, namely the *NcoI* site, made blunt ended by S1 nuclease digestion, and the *HindIII* site, using standard recombinant DNA methodology.

The ends of the KGF cDNA produced by the PCR method were as follows: the 5' end was blunt and began with an ATG codon, followed by the codon TGC for cys residue, number 33, which is the amino terminal residue of the mature form of KGF (see Fig. II-1), and then the entire KGF coding sequence. The stop codon, TAA, and the four bases immediately following, TTGC, were also included on the 3' end of the cDNA. The primer used in the PCR method to direct DNA synthesis to the desired position on the 3' end of the cDNA included a *HindIII* site for insertion of the amplified cDNA into the vector DNA.

Production of antibodies against KGF and KGF-related peptides. Monoclonal antibodies were raised in mice against intact, purified protein from human fibroblasts using 5 or more subcutaneous injections. Test bleeds were screened with a solid-phase (ELISA) assay using highly purified KGF from the conditioned medium of human-- fibroblast cells as antigen. Hybridomas were prepared by routine methods and supernatants were screened with the ELISA assay to detect KGF-reactive antibodies. Positive clones were serially subcloned by the usual methods, and selected subclones were grown as ascites tumors in mice for production of large amounts of antibodies. Antibodies were purified from ascites fluids employing standard techniques (e.g., hydroxyapatite or immunoaffinity resins).

Polyclonal antibodies against a synthetic peptide were raised in rabbits by standard methods, as follows. The peptides were made by solid phase technology and coupled to thyroglobulin by reaction with glutaraldehyde. Serial subcutaneous injections were made and test bleeds were screened by ELISA as well as other techniques, including Western blot analysis and mitogenesis bioassay. IgG immunoglobulins were isolated by affinity chromatography using immobilized protein G.

Polyclonal antibodies were raised in rabbits against both naturally secreted KGF from human fibroblasts and recombinant KGF produced in *E. coli* (see next section), using the following protocol:

i) Initial injection and first boost were administered in the inguinal lymph nodes;

ii) subsequent boosts were made intramuscularly.

5 Screening of test bleeds included ELISA as well as Western blot analysis and mitogenesis bioassay, and IgG was purified as for antibodies against synthetic peptides, above.

## RESULTS

Isolation of cDNA clones encoding the novel growth factor. To search for cDNA clones corresponding to the KGF coding sequence, two pools of oligonucleotides with  
5 lengths of 26 bases were generated based upon a nine-amino acid sequence, Asn-Asp-Met-Thr-Pro-Glu-Gln-Met-Ala, as determined by microsequencing of purified KGF (see Experimental Section I, above and reference II-3. One oligonucleotide pool contained a mixture of all 256  
10 possible coding sequences for the nine amino acids, while the other contained inosine residues at the degenerate third position of the codons for Thr and Pro.

This latter design reduced the number of possible coding sequences in the pool to 16. Inosine in a tRNA  
15 anticodon can form hydrogen bonds with A, C or U (Crick, F.H.C. (1966) *J. Mol. Biol.* 19, 548-555), and oligonucleotides that contain deoxyinosine have been shown to hybridize efficiently with the corresponding cDNA (Ohtsuka, E., Matsuki, S., Ikehara, M., Takashi, Y. and  
20 Matsubara, K. (1985) *J. Biol. Chem.* 260, 2605-2608).

A cDNA library was constructed in a cDNA cloning vector, pCEV9 (Miki, T., Matsui, T., Heidaran, M. and Aaronson, S.A., unpublished observations) using polyadenylated RNA extracted from the human embryonic lung  
25 fibroblast cell line M426 (Aaronson, S.A. and Todaro, G.J. 1968, *Virology* 36, 254-261), the initial source of the growth factor. Screening of the library ( $9 \times 10^5$  plaques) with the  $^{32}\text{P}$ -labelled 26-mer oligonucleotides identified 88 plaques which hybridized to both pools of  
30 oligonucleotide probes.

Characterization and sequencing of selected cDNA clones. Of 10 plaque-purified clones that were analyzed, one, designated clone 49, had a cDNA insert of 3.5 kb, while the rest had inserts ranging from 1.8 kb to 2.1 kb.

Analysis of the smaller clones revealed several common restriction sites. Sequencing of a representative smaller clone, designated clone 32, along with clone 49, demonstrated that they were overlapping cDNAs (Fig II-1A).

Description of the sequence encoding the KGF polypeptide. Alignment of the two cDNAs (clones 32 and 49) established a continuous sequence of 3.85 kb containing the complete KGF coding sequence (Fig. II-1B). An ATG likely to be an initiation codon was located at nucleotide position 446, establishing a 582-base pair open reading frame that ended at a TAA termination codon at position 1030. This open reading frame would encode a 194-amino acid polypeptide with a calculated molecular weight of 22,512 daltons.

The sequence flanking the ATG codon did not conform to the proposed GCC(G/A)CCATGG consensus for optimal initiation by eukaryotic ribosomes (Kozak, M. (1987) *Nucl. Acids Res.* 13, 8125-8148), however, there was an A three nucleotides upstream of the ATG codon. An A at this position is the most highly conserved nucleotide in the consensus. This ATG codon was preceded 85 nucleotides upstream by a TGA stop codon in the same reading frame.

A 19-amino acid sequence that was homologous to the experimentally determined NH<sub>2</sub>-terminus of purified KGF began 32 amino acids downstream of the proposed initiation codon. There was complete agreement between the predicted and experimentally determined amino acid sequences, where unambiguous assignments could be made.

To search for homology between KGF and any known protein, a computer search of the National Biomedical Research Foundation data base using the FASTP program of Lipman and Pearson was conducted (Lipman, D.J. and Pearson, R.W. (1985) *Science* 227, 1435-1443). By this approach, a striking degree of relatedness between the predicted primary structure of KGF and those of acidic and basic FGF, as well as the related *hst*, FGF-5 and *int-2*-encoded proteins was revealed.

Expression of mRNA transcripts of the KGF gene in human cells. In preliminary attempts to examine expression of KGF mRNA in human cells, a probe spanning the majority of the KGF coding sequence (Probe A, Figure II-1A) detected a single 2.4 kb transcript by Northern blot analysis of total M426 RNA (Figure II-1C). This was considerably shorter than the length of the composite cDNA sequence, 3.85 kb.

However, on screening poly(A)-selected M426 RNA, an additional transcript of approximately 5 kb was detected. Furthermore, a probe derived from the untranslated region of clone 49, 3' to the end of clone 32 (Probe B, Figure II-1A), hybridized only to the larger message (Figure II-1C). Thus, it appears that the KGF gene is transcribed as to alternate RNAs. Two other members of the FGF gene family, bFGF (Abraham J.A., et al. (1986) *Science* 233, 545-548) and int-2 (Mansour, S.L. and Martin, G.R. (1988) *EMBO J.* 1, 2035-2041), also express multiple RNAs, the significance of which remains to be determined.

To investigate the normal functional role of KGF, the expression of its transcript in a variety of human cell lines and tissues was examined. As shown in Figure II-3, the predominant 2.4 kb KGF transcript was detected in each of several stromal fibroblast lines derived from epithelial tissues of embryonic, neonatal and adult sources, but not from epithelial cell lines of normal origin. The transcript was also detected in RNA extracted from normal adult kidneys and organs of the gastrointestinal tract, but not from lung or brain. The striking specificity of KGF RNA expression in stromal cells from epithelial tissues indicated that this factor plays a normal role in mesenchymal stimulation of epithelial cell growth.

For comparison, the mRNAs of other growth factors with known activity on epithelial cells were also analyzed in the same tissues as listed above. Among the epithelial and stromal cell lines analyzed, there was no consistent pattern of expression of aFGF or bFGF transcripts (Fig. II-3). The EGF transcript was not

expressed in any of the same cell lines, and was only observed in kidney, among the various tissues. Finally, the TGF $\alpha$  message was not detected in any of the stromal fibroblast lines and was expressed at varying levels in each of the epithelial cell lines. It was also detected at low levels in kidney among the tissues examined (Fig. II-3).

Inhibition of KGF mitogenic activity by heparin.

Heparin has been shown to substantially increase the mitogenic activity of aFGF for a variety of target cells in culture, and to stabilize it from heat inactivation (II-21 and II-22). Despite binding tightly to bFGF, heparin had minimal effects on its mitogenic activity II-22. In view of the relatedness of KGF to the FGFs, the effect of heparin on KGF mitogenic activity was examined. As shown in Table II-1, thymidine incorporation by BALB/MK cells in response to KGF was inhibited 16 fold when heparin was included in the culture medium. In contrast, the activities of both aFGF and bFGF were increased by the same treatment.

Production of anti-KGF antibodies. Several kinds of antibodies which recognize KGF or KGF-like polypeptides have been prepared using standard methodologies well known in the art of experimental immunology and summarized in the Methods section, above. There include: monoclonal antibodies raised in mice against intact, purified protein from human fibroblasts; polyclonal antibodies raised in rabbits against synthetic peptides with sequences based on amino acid sequences predicted from the KGF cDNA sequence; polyclonal antibodies raised in rabbits against both naturally secreted KGF from human fibroblasts and recombinant KGF produced in *E. coli* (see next section).

Monoclonal antibodies from three different hybridomas have been purified. All three recognize the recombinant

as well as the naturally occurring KGF in a solid-phase (ELISA) assay. None cross-reacts with KGF under denaturing conditions (in a Western blot), and none neutralizes mitogenic activity of KGF in the BALB/MK bioassay.

Polyclonal antibodies were generated with a synthetic peptide with the amino acid sequence NDMTPEQMATNVR, corresponding to residues numbered 32 through 44 in KGF (see Fig. II-1), plus an R (Arg) residue instead of the actual Asn residue encoded by the cDNA at position 45. The Asn residue is probably glycosylated in the natural KGF polypeptide and, therefore, was not identified in the amino acid sequencing data obtained directly from that polypeptide (see Discussion, below). Polyclonal antibodies generated with this synthetic peptide recognize both naturally occurring and recombinant KGF in ELISA and Western blot analyses at a level of sensitivity of at least as low as 10 ng protein. These antibodies, however, do not neutralize mitogenic activity of KGF in the BALB/MK bioassay.

Polyclonal antisera against intact natural KGF protein recognizes KGF in both ELISA and Western blot assays. Such antibodies also appear to inhibit mitogenic activity of KGF in the BALB/MK bioassay.

Expression of KGF cDNA in *E. coli*. KGF cDNA was expressed to produce polypeptide, in *E. coli* by placing its coding sequence under control of the hybrid trk promoter (comprising elements of trp and lac promoters), in the plasmid pKK233-2 (Amman, E. and Brosius, J. (1985) Gene 40, 183). To accomplish this, a specific length of KGF cDNA that contained the information to code for the mature KGF molecule (i.e., without its signal peptide) was amplified using the polymerase chain reaction technique (Sakai, R.K., Scharf, S., Faloona, F., Mullis, K.B., Norn, G.T., Erlich, H.A. and Arnheim, N. (1985) Science 230, 1350-1354). The fragment was directionally inserted between two sites in the vector, namely the *Nco*I site, made blunt ended by S1 nuclease digestion, and the *Hind*III site, using standard recombinant DNA methodology.



Selected recombinants were sequenced at their cDNA 5' ends to ensure correct alignment of the ATG initiation codon with the regulatory elements of the trk promoter.

5        Several recombinants were tested for protein production by the usual small scale methods. In brief, the clones were grown to mid-exponential phase ( $OD_{595} \sim 0.5$ ), treated with 1 mM isopropyl  $\beta$ -D-thiogalactopyranoside (IPTG) for 90 minutes, and cell extracts were run on SDS-polyacrylamide gels for Western blot analysis.

10      All recombinants tested synthesized a protein that was recognized by antibodies raised against an amino-terminal KGF peptide. One recombinant was selected which showed the greatest induction from IPTG, for further protein analyses.

15        One liter of bacteria was grown up in NZY broth containing 50  $\mu$ g/ml ampicillin and 12.5  $\mu$ g/ml tetracycline, to  $OD_{595} \sim 0.5$ , and treated for 90 min. with IPTG. The cells were collected by centrifugation, resuspended in 50 mM sodium phosphate (pH 7.3), 0.2 M NaCl, and lysed

20      by sonication. Cell debris was removed by centrifugation, and lysate applied directly to a heparin-Sepharose affinity column.

As determined by Western blot analysis and mitogenic activity in keratinocytes, recombinant KGF was eluted in

25      0.5-0.6 M NaCl. Subsequent purification of the HSAC material with a Mono-S (FPLC) column (Pharmacia) yielded a preparation of KGF estimated to be  $\geq 90\%$  pure, as judged by electrophoretic analysis using SDS-polyacrylamide gels and silver-staining.

30        Recombinant KGF efficiently stimulated thymidine incorporation into BALB/MK keratinocyte cells, but was only marginally active on NIH/3T3 fibroblasts. Half-maximal stimulation of the BALB/MK cells in the standard keratinocyte bioassay was achieved with a concentration

35      of between 2 to 5 ng/ml, compared to a concentration of 10 to 15 ng/ml for KGF purified from M426 cells. One liter of bacterial cells yielded approximately 50  $\mu$ g of Mono-S purified recombinant KGF.

Construction of a chimera containing KGF and aFGF sequences. The studies above indicated that KGF possessed two distinctive characteristics which might be encoded by distinct portions or domains of the polypeptide sequence, as is well known to occur in coding sequences of other multifunctional polypeptides. To test this possibility, a chimeric DNA segment encoding the NH<sub>2</sub>-terminal sequence of KGF grafted onto the C-terminal core of aFGF was constructed, as follows. A *SauI* restriction enzyme site (CCTGAGG) in the 5' end of the KGF cDNA, within codons for residues 76, 77, and 78 (Tyr, Leu, and Arg, respectively; see Fig. II-1) was cut and joined to an homologous site in the aFGF cDNA within codons for amino acids 37 (Phe), 38 (Leu) and 39 (Arg). The 3' and 5' ends of this chimeric DNA were joined to the vector DNA of the plasmid pKK233-2 by the same method used for insertion of the KGF cDNA encoding the secreted form of polypeptide (see Methods, above).

When recombinant *E. coli* cells were constructed using the vector carrying the chimera, and expression tests were conducted as described for mature KGF, above, a novel product with properties of both KGF and aFGF was produced. The peptide was enriched by heparin-Sepharose chromatography and found to have a target cell preference for keratinocytes, like KGF, with minimal activity on fibroblasts (NIH/3T3). The mitogenic activity of this chimeric polypeptide lacks, however, susceptibility to inhibition by heparin, a characteristic which parallels that of aFGF rather than KGF. In fact, the mitogenic activity on keratinocytes is actually enhanced by heparin, as is the case for aFGF. Thus the peptide domains responsible for target cell specificity and heparin sensitivity are clearly distinct and readily separable in KGF, according to the practice of the present invention.

#### DISCUSSION

The experiments described in this section illustrate the practice of several principal embodiments of the present invention. These include isolation of cDNAs encoding KGF, expression of such cDNAs in recombinant cells, production of various antibodies reactive with KGF, and construction and expression of a chimeric cDNA encoding a novel growth factor with amino acid sequences and related functionalities of both KGF and aFGF. The following points related to these embodiments may also be noted to enhance the understanding of the present invention.

The sequence predicted from the KGF cDNA agreed with the amino acid sequence determined from the purified KGF form secreted by human fibroblasts. Moreover, the sequence offered potential explanations for positions where definitive amino acid assignments could not be made by direct amino acid sequencing. Residues 32 and 46 are predicted from the cDNA sequence to be cysteines, and hydrolyzed derivatives of unmodified cysteine residues are not detectable following Edman degradation. The predicted KGF amino acid sequence also contained one potential N-linked glycosylation site (Asn-X-Ser/Thr) from residues 45 through 47. If Asn 45 were glycosylated, it would not be detected by the amino acid sequencing methods employed here. In fact, KGF migrates as a broad band on NaDodSO<sub>4</sub>/PAGE at a higher molecular weight than predicted for the purified protein. This may be accounted for by glycosylation.

The FGFs are heparin-binding mitogens with broad target cell specificities (Thomas, K. (1987) *FASEB J.* 1, 434-440). FGF-5 and *hst* are transforming genes originally detected by DNA-mediated gene transfer. The product of the *int-2* gene is expressed normally during mouse embryogenesis (Jakobovits, A., Shackleford, G.M., Varmus, H.E. and Martin, G.R. (1986) *Proc. Natl. Acad. Sci. USA* 83, 7806-7810) and aberrantly after proviral integration of mouse mammary tumor virus (Peters, G., Brookes, S. and Dickson, S. (1983) *Cell* 33, 364-377). The *hst* gene was identified as a transforming gene from a human stomach tumor (II-11), adjacent normal stomach tissue (II-12) and from Kaposi's sarcoma (II-13), by standard NIH/3T3 transfection assays.

KGF is the sixth member of the fibroblast growth factor family to be identified (Zhan, X., Bates, B., Hu, X. and Goldfarb, M. (1988) *Mol. Cell. Biol.* 8, 3487-3495). While the name FGF-6 does not seem suitable because KGF is devoid of activity on fibroblasts, this nomenclature may also be used for this growth factor, to denote its structural relationship to the FGF family. As all previously characterized growth factors either exclude epithelial cells as targets or include them among a number of sensitive target cells, the highly specific nature of KGF mitogenic activity for epithelial cells, and the sensitivity of keratinocytes in particular, make it unique.

In studies to date, expression of the KGF transcript appears to be specific for stromal cells derived from epithelial tissues, suggesting its function in normal epithelial cell proliferation. The availability of the KGF cDNA clone will make it possible to determine whether abnormal expression of this growth factor can be implicated in clinical conditions characterized by epithelial cell dysplasia and/or neoplasia. Moreover, the ability to produce large quantities of this novel growth factor by recombinant techniques should allow testing of its clinical applicability in situations where specific growth of epithelial cells is of particular importance.

Alignment of the KGF sequence with the five other proteins of the FGF family revealed two major regions of homology, spanning amino acids 65-156 and 162-189 in the predicted KGF sequence, which were separated by a short, nonhomologous series of amino acids with varying lengths in different members of the family (Fig. II-2). In the case of *int-2*, the length of this sequence was 17 residues, while in *hst*, the two homologous regions were contiguous. In KGF the intervening sequence consisted of five amino acids.

In the aligned regions, the KGF amino acid sequence was about 44% identical to *int-2* (mouse), 41% identical to FGF-5 (human), 39% identical to bFGF (human), 37% identical to aFGF (human) and 33% identical to *hst*

(human). In this same region, all six proteins were identical at 19% of the residues, and allowing for conservative substitutions, they showed 28% homology.

5 As shown in Fig. II-2, the amino termini of these related proteins are nonhomologous and of variable length. The primary KGF, FGF-5 and *hst* translation products contain hydrophobic N-terminal regions which likely serve as signal sequences (von Heijne, G. (1986) *Nucl. Acids Res.* 14, 4683-4690). The fact that this N-  
10 terminal domain is not present in the mature KGF molecule (Fig. II-1B) further supports this conclusion. In contrast, the FGFs are synthesized apparently without signal peptides (Thomas, K. (1987) *FASEB J.* 1, 434-440). The *int-2* protein contains an atypically short region of  
15 N-terminal hydrophobic residues (von Heijne, G. (1986) *Nucl. Acids Res.* 14, 4683-4690), but it is not known if the protein is secreted. Moreover, the *int-2* protein contains a long C- terminal extension compared to the other family members.

20 Purified KGF contains five cysteine residues, two of which are conserved throughout the family of FGF related proteins (Fig. II-2). Also of note are the five pairs of basic residues throughout the KGF sequence. This same pattern has been observed in other FGF family members and  
25 may be involved in their interaction with heparin (Schwarzbauer, J.E., Tamkum, J.M., Lemischka, I.R. and Hynes, R.O. (1983) *Cell* 35, 421-431). Dibasic sites are also common targets for proteolytic processing and such processing might account for the microheterogeneity  
30 observed in some KGF preparations (unpublished data).

The KGF cDNA sequence was AT rich throughout its length, but particularly so in the 3' untranslated region where the AT content was 70% as compared to 60% in the putative coding sequence and 63% in the 5' untranslated  
35 region. The 3' untranslated region contained a large number of ATTTA sequences, which have been proposed to be involved in the selected degradation of transiently expressed, unstable RNAs (Shaw, G. and Kamen, R. (1986) *Cell* 46, 659-667). There was no classical AATAAA

polyadenylation signal but two variant sequences, AATTAA and AATACA (Birnstiel, M.L., Busslinger, M. and Strub, K. (1985) *Cell* 41, 349-359), were detected 24 and 19 nucleotides, respectively, upstream of the poly(A) sequence at the 3' end of the cDNA.

It has been suggested that the heparin effect on acidic FGF is either due to stabilization of the active conformation of the growth factor or to formation of a tertiary complex with acidic FGF and its receptor (Schrieber, A.B., Kenny, J., Kowalski, W., Friesel, J., Mehlman, T. and Maciag, T. (1985) *Proc. Natl. Acad. Sci. USA* 82, 6138-6142, Gospodarowicz, O. and Cheng, J. (1986) *J. Cell Physiol.* 128, 475-485). If so, heparin may stabilize a conformation of KGF that is not as active as the free molecule, or form a tight complex that is unable to efficiently interact with its receptor.

While its ability to bind heparin reflects the structural similarities of KGF with the FGF's, the differences in target cell specificities between these related mitogens is remarkable. The FGF's induce division of most nonterminally differentiated cells of both embryonic mesodermal and neuroectodermal origin. In addition to fibroblasts and vascular endothelial tissues, mesodermally derived targets in culture include myoblasts, chondrocytes and osteoblasts (Thomas, K.A. and Gimenez-Gallego, G. (1986) *Trends Biochem. Soc.* 11, 81-84). FGF's are also mitogenic for glial astrocytes and neuroblasts (Gensburger, C., Labourdette, G. and Sensembrenner, M. (1987) *FEBS Lett.* 217, 1-5). The product of the oncogene isolated from Kaposi's sarcoma, which is identical to *hst*, also stimulates proliferation of NIH/3T3 and capillary endothelial cells (Delli-Bovi, P., Curatola, A.M., Kern, F.G., Greco, A., Ittman, M. and Basilico, C. (1987) *Cell* 50, 729-737). To date, KGF induced mitogenesis has only been observed in epithelial cells, and the absence of any detectable activity in fibroblasts or endothelial cells has also been demonstrated (see Experimental Section I, above and II-3

It seems likely, therefore, that KGF acts through a different cell surface receptor than the FGFs.

There is no significant N-terminal homology between KGF and other FGF-related proteins. Thus, the construction of chimeric molecules between KGF and a prototype FGF was undertaken to determine whether the KGF N-terminal domain is sufficient to account for its unique target cell specificity. The results on the first such recombinant polypeptide sequence indicate that the N-terminal domain of KGF essentially encodes the cell preference for keratinocytes, while the susceptibility of KGF to heparin is encoded somewhere in the C-terminal core region which was replaced by sequences of aFGF. This novel KGF-like growth factor may have advantages in clinical applications where administration of an epithelial-specific growth factor is desirable in the presence of heparin, a commonly used anticoagulant. Additional studies on chimeras should also provide insights into which specific domains in the C-terminal core contribute the different effects of heparin on their biologic activities.

For purposes of completing the background description and present disclosure, each of the published articles, patents and patent applications heretofore identified in this specification is hereby incorporated by reference into the specification.

The foregoing invention has been described in some detail for purposes of clarity and understanding. It will also be obvious that various combinations in form and detail can be made without departing from the scope of the invention.

WHAT IS CLAIMED IS:

1. A human keratinocyte growth factor (KGF) having an apparent molecular weight of about 28 kDa as determined by migration in NaDodSO<sub>4</sub>/PAGE, and a specific activity of at least about  $3.4 \times 10^4$  units per milligram of protein, where one unit of activity is defined as that amount which causes half of the maximal possible stimulation of DNA synthesis in BALB/MK keratinocyte cells under standard assay conditions.

2. Human KGF according to claim 1, wherein said specific activity is at least about  $3.2 \times 10^5$  units per milligram protein.

3. A bioassay for KGF-like activity in a test sample which comprises the following steps:

- i) growing keratinocytes in culture to confluence and maintaining said confluent culture in serum-free medium;
- ii) adding a test sample to said confluent culture of keratinocytes;

and

- iii) determining the stimulation of DNA synthesis in said keratinocytes.

4. A method of producing KGF from cultured cells comprising the following steps:

- i) Culturing KGF-producing cells in culture medium under conditions such that KGF is produced;
- ii) concentrating said culture medium so that a first concentrate is formed;
- iii) contacting said concentrate with heparin under conditions such that KGF present in said first concentrate binds to the heparin whereby a heparin-KGF complex is formed;



- iv) separating said heparin-KGF complex from said concentrate;
- v) treating said heparin-KGF complex under conditions such that said KGF dissociates from the heparin so that a solution of free KGF is formed;
- vi) concentrating said solution so that a second concentrate is formed;
- vii) fractionating said second concentrate so that KGF is separated from the remaining components.

5. A method of producing KGF from cultured cells, according to claim 4, wherein said KGF-producing cells are M426 human embryonic fibroblasts.

6. A DNA segment encoding a human keratinocyte growth factor (KGF) protein.

7. A DNA segment, according to claim 6, wherein said protein has the amino acid sequence defined in Figure II-1.

8. A DNA segment encoding a chimeric KGF-like protein which comprises within a single polypeptide molecule functional segments of human KGF and at least one other polypeptide of the fibroblast growth factor family.

9. A recombinant DNA molecule comprising a DNA segment according to claim 6 or claim 8 and a vector.

10. A culture of cells transformed with said recombinant DNA molecule according to claim 9.

11. A method of producing a human KGF protein comprising culturing said cells according to claim 10 in a culture medium under conditions such that said protein is produced and isolating said protein from said cells.

12. A method of producing a human KGF protein comprising culture said cells according to claim 10 in a culture medium, wherein said protein is secreted from said cell, and isolating said protein from said medium.

13. A human KGF or KGF-like protein having the amino acid sequence in Figure II-1B.

14. A human KGF or KGF-like protein, according to claim 13, which is not glycosylated.

15. An antibody specific for a peptide having the amino acid sequence of human KGF or KGF-like protein, according to claim 13.

16. The antibody according to claim 15 which neutralizes the mitogenic activity of human KGF.

17. A bioassay for expression of a gene encoding KGF, comprising the steps of:

- i) isolating mRNA from tissues or cells;
- and
- ii) annealing said RNA to a DNA probe encoding a human KGF;
- iii) determining the amount of DNA:RNA hybrid containing said DNA probe.

18. A bioassay for KGF antigen comprising the steps of:

- i) extracting polypeptides from body fluids or tissue samples;
- and
- ii) determining the level of human KGF antigen by reaction with an antibody specific for a peptide having the amino acid sequence of human KGF or KGF-like protein, according to claim 13.

19. A pharmaceutical composition for treatment of conditions requiring specific stimulation of epithelial cells, comprising KGF according to claim 1 or claim 13, and an acceptable pharmaceutical carrier.

20. A pharmaceutical composition for treatment of conditions requiring specific inhibition of stimulation of epithelial cells by KGF, comprising antibodies to KGF according to claim 15, and an acceptable pharmaceutical carrier.

ABSTRACT

Discoveries are disclosed that show particular aspects of recombinant DNA technology can be used successfully to produce hitherto unknown human keratinocyte growth factor (KGF) protein free of other polypeptides. These proteins can be produced in various functional forms from spontaneously secreting cells or from DNA segments introduced into cells. These forms variously enable biochemical and functional studies of this novel protein as well as production of antibodies. Means are described for determining the level of expression of genes for the KGF protein, for example, by measuring mRNA levels in cells or by measuring antigen secreted in extracellular or body fluids.

Table I-1. Growth Factor Purification

Purification step	Protein (mg)	Total activity (units)*	Specific activity (units/mg)
Conditioned medium (10 liters)	$1.4 \times 10^{3a}$	$2.5 \times 10^4$	$1.8 \times 10^1$
Ultrafiltration (retentate)	$1.3 \times 10^{3a}$	$3.2 \times 10^4$	$2.5 \times 10^1$
HSAC	0.73 <sup>b</sup>	$1.6 \times 10^4$	$2.2 \times 10^1$
0.6 MM NaCl pool			
TSK-G3000 SW	$8.4 \times 10^{-3b}$	$2.7 \times 10^3$	$3.2 \times 10^5$
C <sub>4</sub> -HPLC	$6.1 \times 10^{-3b}$	$2.1 \times 10^2$	$3.4 \times 10^4$

Recoveries were calculated by assuming that all of the mitogenic activity in the starting material was due to the isolated factor.

5 \*One unit of activity is defined as half of the maximal stimulation of thymidine incorporation induced by TSK-purified factor in the BALB/MK bioassay, in which approximately 3 ng of the TSK-purified factor stimulated 1 unit of activity.

10 <sup>a</sup> Protein was estimated by using the Bradford reagent from BioRad.

<sup>b</sup> Protein was estimated by using  $A_{214}^{1\%} = 140$ .

Table I-2 Target Cell Specificity of Growth Factors

Growth Factor	<u>Epithelial</u>			<u>Fibroblast</u>	<u>Endothelial</u>
	BALK/MK	BS/589	CCL208	NIH/3T3S	Human saphenous vein
KGF	500-1000	2-3	5-10	<1	<1
EGF	100-200	20-40	10-30	10-20	n.d.
TGFa	150-300	n.d.	n.d.	10-20	n.d.
aFGF*	300-500	2-3	5-10	50-70	5
bFGF	100-200	2-3	2-5	50-70	5

Comparison of maximal thymidine incorporation stimulated by KGF and other growth factors in a variety of cell lines, expressed as fold stimulation over background.

This data represents a summary of four different experiments.

\*Maximal stimulation by aFGF required the presence of heparin (Sigma), 20 $\mu$ g/ml.

n.d. = not determined.




Figure I-1. Heparin-Sepharose affinity chromatography of conditioned medium from M426 human embryonic fibroblasts. Approximately 150 ml of ultrafiltration retentate derived from five liters of M426 conditioned medium were loaded onto a heparin-Sepharose column (6 ml bed volume) in 1 hr. After washing the column with 150 ml of the equilibration buffer, 20 mM Tris-HCl, pH 7.50/0.3M NaCl, the retained protein (<5% of the total protein in the retentate) was eluted with a modified linear gradient of increasing NaCl concentration. Fraction size was 3.8 ml and flow rate during gradient elution was 108 ml/hr. Two  $\mu$ l of the indicated fractions were transferred to microtiter wells containing a final volume of 0.2 ml for assay of <sup>3</sup>H-thymidine incorporation in BALB/MK cells as described in the Methods.

Figure I-2. SEE LEGEND NEXT PAGE






Figure I-2. (A) Reversed-phase  $C_4$ HPLC of BALB/MK mitogenic activity. Active fractions eluted from heparin-Sepharose with 0.6M NaCl were processed with the Centricon -10 and loaded directly onto a  $C_4$  Vydac column (4.6 x 250 mm) which had been equilibrated in 0.1% trifluoroacetic acid/20% acetonitrile (ACN). After washing the column with 4 ml of equilibration buffer, the sample was eluted with a modified linear gradient of increasing % ACN. Fraction size was 0.2 ml and flow rate was 0.5 ml/min. Aliquots for the assay of  $^3$ H-thymidine incorporation in BALB/MK cells were promptly diluted 10-fold with 50  $\mu$ g/ml bovine serum albumin/20 mM Tris-HCl, pH 7.5, and tested at a final dilution of 200-fold. (B) NaDodSO<sub>4</sub>/PAGE analysis of selected fractions from the  $C_4$  chromatography shown in-panel A. Half of each fraction was dried, redissolved in NaDodSO<sub>4</sub>/2-mercaptoethanol, heat denatured and electrophoresed in a 14% polyacrylamide gel which was subsequently stained with silver. The position of each molecular weight marker (mass in kDa) is indicated by an arrow. (C) DNA synthesis in BALB/MK cells triggered by the fractions analyzed in Panel B. Activity is expressed as the fold stimulation over background which was 100 cpm.




Figure I-3. Molecular sieving HPLC (TSK 3000SW) chromatography of the BALB/MK mitogenic activity. Approximately 50  $\mu$ l of a Centricon-processed, 0.6M NaCl pool from HSAC were loaded onto a LKB GlasPac TSK G3000SW column (8 x 300 mm), previously equilibrated in 20 mM Tris-HCl, pH 6.8/0.5M NaCl, and eluted as 0.2 ml fractions at a flow rate of 0.4 ml/min. Aliquots of 2  $\mu$ l were transferred to microtiter wells containing a final volume of 0.2 ml for assay of <sup>3</sup>H-thymidine incorporation in BALB/MK cells. The elution positions of molecular weight markers (mass in kDa) were as indicated by the arrows.




Figure I-4. Comparison of BALB/MK DNA synthesis in response to TSK-purified mitogen and other growth factors. Incorporation of  $^3\text{H}$ -thymidine into trichloroacetic acid-insoluble DNA, expressed as fold stimulation over background, was measured as a function of the concentration of the indicated growth factors. Background values with no sample added were 150 cpm. The results represent mean values of two independent experiments. Replicates in each experiment were within 10% of mean values. TSK-purified mitogen,  $\bullet$ ; EGF,  $\Delta$ ; aFGF,  $\square$ ; bFGF,  $\circ$ .




Figure I-5. Comparative growth of BALB/MK cells in a chemically defined medium in response to different combinations of growth factors. Cultures were plated at a density of  $2.5 \times 10^4$  cells per dish on 35 mm Petri dishes precoated with poly-D-lysine/fibronectin in a 1:1 mixture of Eagle's minimal essential medium and Ham's F12 medium supplemented with transferrin,  $\text{Na}_2\text{SeO}_3$ , ethanolamine and the growth factors indicated below. After 10 days, the plates were fixed and stained with Giemsa. Key: a) no growth factor; b) EGF alone; c) insulin alone; d) KGF alone; e) EGF and dialyzed fetal calf serum (final concentration, 10%); f) KGF and EGF; g) KGF and insulin; h) EGF and insulin. Final concentrations of the growth factors were as follows: EGF, 20 ng/ml; insulin, 10  $\mu\text{g}/\text{ml}$ ; and KGF, 40 ng/ml.

Table II-1. Effect of Heparin on KGF Mitogen Activity.

	<u>Growth Factor</u>	<u>BALB/MK</u>		<u>NIH/3T3</u>	
		-	+	-	+
5	KGF	150	9.5	<1	<1
	aFGF	106	259	10.4	68
	bFGF	30	124	45.7	70

Cells were plated in microtiter plates, grown to confluence in serum containing media and then placed in a serum-free medium for 24-72 hr prior to sample addition. Mitogenesis assays were performed as described (see Experimental Section I, above II-3. Where indicated, heparin was included in the culture media at a final concentration of 20  $\mu$ g/ml. The concentration of all the growth factors was 50 ng/ml. The results represent fold stimulation of  $^3$ H-thymidine incorporation in the indicated assay cell in the presence (+) or absence (-) of heparin. Each value represents the mean result from two independent experiments in which each point, in turn, represents the mean value of duplicate analyses.

Figure II-1A. SEE LEGEND FOLLOWING

A.




Figure II-1B. SEE LEGEND FOLLOWING

[illegible]

Figure II-1C. SEE LEGEND FOLLOWING






Figure II-1. Nucleotide sequence and deduced amino acid sequence of KGF cDNA, and identification of KGF gene transcripts. (A) Schematic representation of human KGF cDNA clones. Overlapping pCEV9 clones 32 and 49, used in sequence determination, are shown above a diagram of the complete structure in which untranslated regions are depicted by a line and the coding sequence is boxed. The hatched region denotes sequences of the signal peptide. Selected restriction sites are indicated. (B) KGF cDNA nucleotide and predicted amino acid sequences. Nucleotides are numbered on the left; amino acids are numbered throughout. The N-terminal peptide sequence derived from purified KGF is underlined. The hydrophobic N-terminal domain is italicized. The potential asparagine-linked glycosylation site is overlined. The variant polyadenylation signals, AATTAA and AATACA, close to the 3' end of the RNA, are boxed. (C) Identification of KGF mRNAs by Northern blot analysis. Lanes a and c, poly(A)-selected M426 RNA; lanes b and d, total cellular M426 RNA. Filters were hybridized with a <sup>32</sup>P-labeled 695 bp *Bam*HI/*Bcl*I fragment from clone 32 (Probe A, Fig. II-1A), lanes a and b, or a 541 bp *Apa*I/*Eco*RI fragment from clone 49 (Probe B, Fig. II-1A), lanes c and d.




Figure II-2. Topological comparison of the FGF family of related molecules. The two protein domains that share high homology are shown by shaded boxes. Hatched boxes indicate putative signal peptide sequences. The positions of two conserved cysteine residues (C) are shown.

Figure II-3. SEE LEGEND NEXT PAGE




Figure II-3. Northern blot analysis of KGF mRNA in normal human cell lines and tissues, and comparison with mRNA expression of other growth factors with known activity on epithelial cells. Total cellular RNAs were isolated by cesium trifluoro-acetate gradient centrifugation. 10  $\mu$ g of RNA were denatured and electrophoresed in 1% formaldehyde gels. Following milk alkali denaturation (50 mM NaOH for 30'), RNA was transferred to nitrocellulose filters using 1 M ammonium acetate as a convectant. Filters were hybridized to a <sup>32</sup>P-labelled cDNA probe containing the *Bam*HI/*Bcl*II fragment containing the majority of the KGF coding sequence (A) or similar probes from the other growth factor DNAs. The following human cell types were used: squamous cell carcinomas (A253, A388 and A431); mammary epithelial cells B5/589; immortalized bronchial epithelial cells (S6 and R1); keratinocytes immortalized with Ad12-SV40; primary human keratinocytes; neonatal foreskin fibroblasts, (AG1523) adult skin fibroblasts (501T); and embryonic lung fibroblasts (WI-38 and M426).